10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ADIPOR2 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5105C

Product Information

Application	WB, IF, FC, E
Primary Accession	Q86V24
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB23213
Calculated MW	43884
Antigen Region	45-72

Additional Information

Gene ID	79602
Other Names	Adiponectin receptor protein 2, Progestin and adipoQ receptor family member II, ADIPOR2, PAQR2
Target/Specificity	This ADIPOR2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 45-72 amino acids from the Central region of human ADIPOR2.
Dilution	WB~~1:1000 IF~~1:10~50 FC~~1:25 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ADIPOR2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ADIPOR2 (HGNC:24041)
Function	Receptor for ADIPOQ, an essential hormone secreted by adipocytes that regulates glucose and lipid metabolism (PubMed:12802337, PubMed:25855295). Required for normal body fat and glucose homeostasis.

	ADIPOQ-binding activates a signaling cascade that leads to increased PPARA activity, and ultimately to increased fatty acid oxidation and glucose uptake. Has intermediate affinity for globular and full-length adiponectin. Required for normal revascularization after chronic ischemia caused by severing of blood vessels (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein Note=Localized to the cell membrane and intracellular organelles
Tissue Location	Ubiquitous (PubMed:16044242). Highly expressed in skeletal muscle, liver and placenta (PubMed:12802337). Weakly expressed in brain, heart, colon, spleen, kidney, thymus, small intestine, peripheral blood leukocytes and lung (PubMed:12802337)


Background

The adiponectin receptors, ADIPOR1 (MIM 607945) and ADIPOR2, serve as receptors for globular and full-length adiponectin (MIM 605441) and mediate increased AMPK (see MIM 602739) and PPAR-alpha (PPARA; MIM 170998) ligand activities, as well as fatty acid oxidation and glucose uptake by adiponectin.


References

Ferguson, J.F., et al. Am. J. Clin. Nutr. 91(3):794-801(2010) Halvatsiotis, I., et al. Cardiovasc Diabetol 9, 10 (2010) Yoshihara, K., et al. J Reprod Med 54 (11-12), 669-674 (2009)

Images




Anti-ADIPOR2 Antibody (Center) at 1:1000 dilution + Human liver tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 44 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Confocal immunofluorescent analysis of ADIPOR2 Antibody (Center) (Cat. #AP5105c) with Hela cell followed by Alexa Fluor寨?489-conjugated goat anti-rabbit lgG (green). DAPI was used to stain the cell nuclear (blue).

ADIPOR2 Antibody (Center) (Cat. #AP5105c) flow cytometric analysis of Hela cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary



antibodies were used for the analysis.

Citations

- Contrasting Effects of Adipokines on the Cytokine Production by Primary Human Bronchial Epithelial Cells: Inhibitory Effects of Adiponectin
- Novel phytopeptide osmotin mimics preventive effects of adiponectin on vascular inflammation and atherosclerosis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.