

CLCN4 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51090

Product Information

Application	WB
Primary Accession	P51793
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	84917

Additional Information

Gene ID	1183
Other Names	H(+)/Cl(-) exchange transporter 4, Chloride channel protein 4, CIC-4, Chloride transporter CIC-4, CLCN4
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CLCN4. The exact sequence is proprietary.
Dilution	WB~~ 1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	CLCN4
Function	Strongly outwardly rectifying, electrogenic H(+)/Cl(-)exchanger which mediates the exchange of chloride ions against protons (PubMed: 18063579 , PubMed: 23647072 , PubMed: 25644381 , PubMed: 27550844 , PubMed: 28972156). The CLC channel family contains both chloride channels and proton-coupled anion transporters that exchange chloride or another anion for protons (PubMed: 29845874). The presence of conserved gating glutamate residues is typical for family members that function as antiporters (PubMed: 29845874).
Cellular Location	Early endosome membrane {ECO:0000250 UniProtKB:P51794}; Multi-pass membrane protein. Late endosome membrane; Multi-pass membrane protein. Endoplasmic reticulum membrane; Multi-pass membrane protein. Lysosome membrane; Multi-pass membrane protein. Recycling endosome membrane; Multi-pass membrane protein. Note=Localizes to late endosome membrane, lysosome membrane and recycling endosome membrane in the

presence of CLCN3

Tissue Location

Abundant in skeletal muscle and also detectable in brain and heart

Background

Proton-coupled chloride transporter. Functions as antiport system and exchanges chloride ions against protons.

References

van Slegtenhorst M.A., et al. Hum. Mol. Genet. 3:547-552(1994).


Kawasaki M., et al. Am. J. Physiol. 277:C948-C954(1999).

Rae J.L., et al. Submitted (JUL-1999) to the EMBL/GenBank/DDBJ databases.

Ota T., et al. Nat. Genet. 36:40-45(2004).

Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

Images



Anti-CHML Antibody at 1:1000 dilution + H.skeletal muscle tissue lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 74 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.