

Anti-CLCN6 Antibody

Catalog # AP51091

Product Information

Application	WB, IHC, IF/IC
Primary Accession	P51797
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	97289

Additional Information

Gene ID	1185
Other Names	KIAA0046; Chloride transport protein 6; Chloride channel protein 6; CLC-6
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CLCN6. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	CLCN6 (HGNC:2024)
Synonyms	KIAA0046
Function	Voltage-gated channel mediating the exchange of chloride ions against protons. Functions as antiporter and contributes to the acidification of the late endosome lumen. The CLC channel family contains both chloride channels and proton-coupled anion transporters that exchange chloride or another anion for protons. The presence of conserved gating glutamate residues is typical for family members that function as antiporters.
Cellular Location	Late endosome membrane; Multi-pass membrane protein
Tissue Location	Testis, ovary, small intestine, brain and skeletal muscle. Low level expression in aortic and coronary vascular smooth muscle cells, and aortic endothelial cells. Isoform 3 is only detected in kidney.


References

Brandt S.,et al.FEBS Lett. 377:15-20(1995).
Eggermont J.,et al.Biochem. J. 325:269-276(1997).
Kornak U.,et al.Biochim. Biophys. Acta 1447:100-106(1999).
Nomura N.,et al.DNA Res. 1:223-229(1994).
Nakajima D.,et al.DNA Res. 9:99-106(2002).


Images



Western blot analysis of CLCN6 expression in HCT116 (A), SKOVCA3 (B) whole cell lysates. (Predicted band size: 97 kD; Observed band size: 97 kD)



Immunohistochemical analysis of CLCN6 staining in human kidney cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of CLCN6 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.