

Cyclin M2 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51098

Product Information

ApplicationWB, IP, ICCPrimary AccessionQ9H8M5

Reactivity Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW96623

Additional Information

Gene ID 54805

Other Names Metal transporter CNNM2, Ancient conserved domain-containing protein 2,

Cyclin-M2, CNNM2, ACDP2

Target/Specificity KLH-conjugated synthetic peptide encompassing a sequence within the center

region of human Cyclin M2. The exact sequence is proprietary.

Dilution WB~~1:1000 IP~~N/A ICC~~N/A

Format 0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name CNNM2

Synonyms ACDP2

Function Divalent metal cation transporter. Mediates transport of divalent metal

cations in an order of Mg(2+) > Co(2+) > Mn(2+) > Sr(2+) > Ba(2+) > Cu(2+) >

Fe(2+) (By similarity).

Cellular Location Cell membrane; Multi-pass membrane protein

Tissue Location Widely expressed. Expressed at higher level in brain, kidney and placenta,

while it is weakly expressed in skeletal muscle. In the kidney, it is expressed in

the distal convoluted tubule and the thick ascending limb of Henle loop

Background

Divalent metal cation transporter. Mediates transport of divalent metal cations in an order of Mg(2+) > Co(2+) > Mn(2+) > Sr(2+) > Ba(2+) > Cu(2+) > Fe(2+) (By similarity).

References

Ota T., et al. Nat. Genet. 36:40-45(2004).
Deloukas P., et al. Nature 429:375-381(2004).
Wang C.-Y., et al. Gene 306:37-44(2003).
Stuiver M., et al. Am. J. Hum. Genet. 88:333-343(2011).

Citations

• Human CNNM2 is not a Mg(2+) transporter per se.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.