

GPR159 Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51136

Product Information

Application	WB, IHC-P
Primary Accession	P25106
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41493

Additional Information

Gene ID	57007
Other Names	Atypical chemokine receptor 3, C-X-C chemokine receptor type 7, CXCR-7, Chemokine orphan receptor 1, G-protein coupled receptor 159, G-protein coupled receptor RDC1 homolog, RDC-1, ACKR3, CMKOR1, CXCR7, GPR159, RDC1
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human GPR159. The exact sequence is proprietary.
Dilution	WB~~1:1000 IHC-P~~N/A
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C. Stable for 12 months from date of receipt

Protein Information

Name	ACKR3 (HGNC:23692)
Function	Atypical chemokine receptor that controls chemokine levels and localization via high-affinity chemokine binding that is uncoupled from classic ligand-driven signal transduction cascades, resulting instead in chemokine sequestration, degradation, or transcytosis. Also known as interceptor (internalizing receptor) or chemokine-scavenging receptor or chemokine decoy receptor. Acts as a receptor for chemokines CXCL11 and CXCL12/SDF1 (PubMed: 16107333 , PubMed: 19255243 , PubMed: 19380869 , PubMed: 20161793 , PubMed: 22300987). Chemokine binding does not activate G-protein-mediated signal transduction but instead induces beta-arrestin recruitment, leading to ligand internalization and activation of MAPK signaling pathway (PubMed: 16940167 , PubMed: 18653785 , PubMed: 20018651). Required for regulation of CXCR4 protein levels in migrating interneurons, thereby adapting their chemokine responsiveness (PubMed: 16940167 ,

PubMed:[18653785](#)). In glioma cells, transduces signals via MEK/ERK pathway, mediating resistance to apoptosis. Promotes cell growth and survival (PubMed:[16940167](#), PubMed:[20388803](#)). Not involved in cell migration, adhesion or proliferation of normal hematopoietic progenitors but activated by CXCL11 in malignant hematopoietic cells, leading to phosphorylation of ERK1/2 (MAPK3/MAPK1) and enhanced cell adhesion and migration (PubMed:[17804806](#), PubMed:[18653785](#), PubMed:[19641136](#), PubMed:[20887389](#)). Plays a regulatory role in CXCR4-mediated activation of cell surface integrins by CXCL12 (PubMed:[18653785](#)). Required for heart valve development (PubMed:[17804806](#)). Regulates axon guidance in the oculomotor system through the regulation of CXCL12 levels (PubMed:[31211835](#)). Acts as a receptor for SHLP2, mediating its effects on activation of proopiomelanocortin neurons in the arcuate nucleus of the hypothalamus which leads to suppression of food intake and increased energy expenditure (PubMed:[37468558](#)).

Cellular Location

Cell membrane; Multi-pass membrane protein. Early endosome. Recycling endosome. Note=Predominantly localizes to endocytic vesicles, and upon stimulation by the ligand is internalized via clathrin-coated pits in a beta-arrestin-dependent manner. Once internalized, the ligand dissociates from the receptor, and is targeted to degradation while the receptor is recycled back to the cell membrane.

Tissue Location

Expressed in monocytes, basophils, B-cells, umbilical vein endothelial cells (HUVEC) and B-lymphoblastoid cells. Lower expression detected in CD4+ T-lymphocytes and natural killer cells. In the brain, detected in endothelial cells and capillaries, and in mature neurons of the frontal cortex and hippocampus. Expressed in tubular formation in the kidney. Highly expressed in astroglial tumor endothelial, microglial and glioma cells. Expressed at low levels in normal CD34+ progenitor cells, but at very high levels in several myeloid malignant cell lines. Expressed in breast carcinomas but not in normal breast tissue (at protein level).

Background

Atypical chemokine receptor that controls chemokine levels and localization via high-affinity chemokine binding that is uncoupled from classic ligand-driven signal transduction cascades, resulting instead in chemokine sequestration, degradation, or transcytosis. Also known as interceptor (internalizing receptor) or chemokine-scavenging receptor or chemokine decoy receptor. Acts as a receptor for chemokines CXCL11 and CXCL12/SDF1. Chemokine binding does not activate G-protein-mediated signal transduction but instead induces beta-arrestin recruitment, leading to ligand internalization and activation of MAPK signaling pathway. Required for regulation of CXCR4 protein levels in migrating interneurons, thereby adapting their chemokine responsiveness. In glioma cells, transduces signals via MEK/ERK pathway, mediating resistance to apoptosis. Promotes cell growth and survival. Not involved in cell migration, adhesion or proliferation of normal hematopoietic progenitors but activated by CXCL11 in malignant hematopoietic cells, leading to phosphorylation of ERK1/2 (MAPK3/MAPK1) and enhanced cell adhesion and migration. Plays a regulatory role in CXCR4-mediated activation of cell surface integrins by CXCL12. Required for heart valve development. Acts as coreceptor with CXCR4 for a restricted number of HIV isolates.

References

Sreedharan S.P.,et al.Proc. Natl. Acad. Sci. U.S.A. 88:4986-4990(1991).
Oates E.L.,et al.Submitted (OCT-1996) to the EMBL/GenBank/DDBJ databases.
Bi A.,et al.Submitted (OCT-1997) to the EMBL/GenBank/DDBJ databases.
Martin A.L.,et al.Submitted (JUN-2006) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.