

Cytochrome P450 2A13 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51144

Product Information

Application	WB, IP, ICC, IHC-P
Primary Accession	Q16696
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	56688

Additional Information

Gene ID	1553
Other Names	Cytochrome P450 2A13, CYP11A13, CYP2A13
Dilution	WB~~1:1000 IP~~N/A ICC~~N/A IHC-P~~N/A
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	CYP2A13
Function	Exhibits a coumarin 7-hydroxylase activity. Active in the metabolic activation of hexamethylphosphoramide, N,N-dimethylaniline, 2'-methoxyacetophenone, N-nitrosomethylphenylamine, and the tobacco-specific carcinogen, 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone. Possesses phenacetin O-deethylation activity.
Cellular Location	Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome membrane; Peripheral membrane protein
Tissue Location	Expressed in liver and a number of extrahepatic tissues, including nasal mucosa, lung, trachea, brain, mammary gland, prostate, testis, and uterus, but not in heart, kidney, bone marrow, colon, small intestine, spleen, stomach, thymus, or skeletal muscle

Background

Exhibits a coumarin 7-hydroxylase activity. Active in the metabolic activation of

hexamethylphosphoramide, N,N- dimethylaniline, 2'-methoxyacetophenone, N- nitrosomethylphenylamine, and the tobacco-specific carcinogen, 4- (methylnitrosamino)-1-(3-pyridyl)-1-butanone. Possesses phenacetin O-deethylation activity.

References

- Fernandez-Salguero P.,et al.Am. J. Hum. Genet. 57:651-660(1995).
Su T.,et al.Cancer Res. 60:5074-5079(2000).
Cauffiez C.,et al.Biochem. Biophys. Res. Commun. 317:662-669(2004).
DeVore N.M.,et al.Drug Metab. Dispos. 36:2582-2590(2008).
Sansen S.,et al.Arch. Biochem. Biophys. 464:197-206(2007).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.