



# Cytochrome P450 2U1 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51148

#### **Product Information**

**Application** WB, ICC, IHC-P

Primary Accession

Reactivity
Human

Host
Clonality
Polyclonal
Calculated MW

O7Z449
Human
Puman
Rabbit
Polyclonal
61987

#### **Additional Information**

**Gene ID** 113612

Other Names Cytochrome P450 2U1, CYP2U1

**Dilution** WB~~1:1000 ICC~~N/A IHC-P~~N/A

Format 0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

**Storage** Store at -20 °C.Stable for 12 months from date of receipt

### **Protein Information**

Name CYP2U1 {ECO:0000303 | PubMed:14660610,

ECO:0000312 | HGNC:HGNC:20582}

**Function** A cytochrome P450 monooxygenase involved in the metabolism of

arachidonic acid and its conjugates (PubMed:<u>14660610</u>, PubMed:<u>24563460</u>). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons

provided by NADPH via cytochrome P450 reductase (CPR; NADPH-

ferrihemoprotein reductase) (PubMed:<u>14660610</u>, PubMed:<u>24563460</u>). Acts as an omega and omega-1 hydroxylase for arachidonic acid and possibly for other long chain fatty acids. May modulate the arachidonic acid signaling

pathway and play a role in other fatty acid signaling processes

(PubMed: 14660610, PubMed: 24563460). May down-regulate the biological activities of N-arachidonoyl-serotonin, an endocannabinoid that has

anti-nociceptive effects through inhibition of fatty acid amide hydrolase FAAH, TRPV1 receptor and T-type calcium channels. Catalyzes C-2 oxidation of the indole ring of N-arachidonoyl-serotonin forming a less active product

2-oxo-N-arachidonoyl-serotonin (PubMed:24563460).

**Cellular Location** Endoplasmic reticulum membrane; Multi-pass membrane protein. Microsome

membrane; Multi- pass membrane protein. Mitochondrion inner membrane;

Multi-pass membrane protein

**Tissue Location** 

Widely expressed with stronger expression in thymus, heart and cerebellum.

# **Background**

Catalyzes the hydroxylation of arachidonic acid, docosahexaenoic acid and other long chain fatty acids. May modulate the arachidonic acid signaling pathway and play a role in other fatty acid signaling processes.

## References

Chuang S.S.,et al.J. Biol. Chem. 279:6305-6314(2004). Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases. Karlgren M.,et al.Biochem. Biophys. Res. Commun. 315:679-685(2004). Choudhary D.,et al.Arch. Biochem. Biophys. 436:50-61(2005). Tesson C.,et al.Am. J. Hum. Genet. 91:1051-1064(2012).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.