

Anti-Cytochrome P450 4Z1/2 Antibody

Catalog # AP51153

Product Information


Application	WB, IHC, IF/IC
Primary Accession	Q5VVE4
Other Accession	Q8N1L4
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59 KDa

Additional Information

Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Cytochrome P450 4Z1/2. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Images




Western blot analysis of Cytochrome P450 4Z1/2 expression in PC12 (A), MEF (B), HeLa (C), MCF7 (D) whole cell lysates. (Predicted band size: 40 kD; Observed band size: 59; 40 kD)

Immunohistochemical analysis of Cytochrome P450 4Z1/2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was



pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of Cytochrome P450 4Z1/2 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.