

E2F4 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51173

Product Information

| Application | WB, IHC-P |
|-------------------|---------------|
| Primary Accession | <u>Q16254</u> |
| Reactivity | Human, Mouse |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 43960 |

Additional Information

| Gene ID | 1874 |
|-------------|-----------------------------------------------------------|
| Other Names | Transcription factor E2F4, E2F-4, E2F4 |
| Dilution | WB~~1:1000 IHC-P~~N/A |
| Format | 0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50% |
| Storage | Store at -20 °C.Stable for 12 months from date of receipt |

Protein Information

| Name | E2F4 |
|-------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Function | Transcription activator that binds DNA cooperatively with DP proteins through the E2 recognition site, 5'-TTTC[CG]CGC-3' found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication. The DRTF1/E2F complex functions in the control of cell-cycle progression from G1 to S phase. E2F4 binds with high affinity to RBL1 and RBL2. In some instances can also bind RB1. Specifically required for multiciliate cell differentiation: together with MCIDAS and E2F5, binds and activate genes required for centriole biogenesis. |
| Cellular Location | Nucleus. |
| Tissue Location | Found in all tissue examined including heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas |

Background

Transcription activator that binds DNA cooperatively with DP proteins through the E2 recognition site,

5'-TTTC[CG]CGC- 3' found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication. The DRTF1/E2F complex functions in the control of cell-cycle progression from G1 to S phase. E2F4 binds with high affinity to RBL1 and RBL2. In some instances can also bind RB1.

References

Ginsberg D.,et al.Genes Dev. 8:2665-2679(1994). Beijersbergen R.L.,et al.Genes Dev. 8:2680-2690(1994). Sardet C.,et al.Proc. Natl. Acad. Sci. U.S.A. 92:2403-2407(1995). Schwemmle S.,et al.Int. J. Cancer 86:672-677(2000). Goshima N.,et al.Nat. Methods 5:1011-1017(2008).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.