10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-ELMO1 Antibody

Catalog # AP51187

Product Information

Application	WB, IF/IC
Primary Accession	Q92556
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	83829

Additional Information

Gene ID	9844
Other Names	KIAA0281; Engulfment and cell motility protein 1; Protein ced-12 homolog
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human ELMO1. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	ELMO1
Synonyms	KIAA0281
Function	Regulates the activity of DOCK guanine nucleotide exchange factors (GEFs) (PubMed:11595183, PubMed:12134158, PubMed:32651375, PubMed:38857861). Acts in association with DOCK1, DOCK2, DOCK5 and CRK (PubMed:11595183, PubMed:12134158, PubMed:32651375, PubMed:38857861). Promotes RAC1 activation by DOCK5; the activity is stimulated by RHOG (PubMed:38857861). Involved in cytoskeletal rearrangements required for phagocytosis of apoptotic cells and cell motility (PubMed:11595183).
Cellular Location	Cytoplasm. Cell membrane. Note=Translocation to plasma membrane seems to be mediated by DOCK1 and CRK
Tissue Location	Widely expressed, with a higher expression in the spleen and placenta


References

Gumienny T.L.,et al.Cell 107:27-41(2001).
Ohara O.,et al.DNA Res. 4:53-59(1997).
Wiemann S.,et al.Genome Res. 11:422-435(2001).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

Images



Western blot analysis of ELMO1 expression in mouse liver (A), mouse brain (B), rat liver (C), rat brain (D) whole cell lysates. (Predicted band size: 83 kD; Observed band size: 90 kD)



Immunofluorescent analysis of ELMO1 staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.