

Anti-GLP-1 Receptor Antibody

Catalog # AP51232

Product Information

Application	WB, IHC, IF/IC
Primary Accession	P43220
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53026

Additional Information

Gene ID	2740
Other Names	Glucagon-like peptide 1 receptor; GLP-1 receptor; GLP-1-R; GLP-1R
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human GLP-1 Receptor. The exact sequence is proprietary.
Dilution	WB~~ 1:1000 IHC~~1:100~500 IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	GLP1R
Function	G protein-coupled receptor for glucagon-like peptide 1 (GLP- 1) (PubMed: 19861722 , PubMed: 26308095 , PubMed: 27196125 , PubMed: 28514449 , PubMed: 7517895 , PubMed: 8216285 , PubMed: 8405712). Ligand binding triggers activation of a signaling cascade that leads to the activation of adenylyl cyclase and increased intracellular cAMP levels (PubMed: 19861722 , PubMed: 26308095 , PubMed: 27196125 , PubMed: 28514449 , PubMed: 7517895 , PubMed: 8216285 , PubMed: 8405712). Plays a role in regulating insulin secretion in response to GLP-1 (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein

References


Thorens B.,et al.Diabetes 42:1678-1682(1993).

Dillon J.S.,et al.Endocrinology 133:1907-1910(1993).
 Graziano M.P.,et al.Biochem. Biophys. Res. Commun. 196:141-146(1993).
 van Eyll B.,et al.FEBS Lett. 348:7-13(1994).
 Wei Y.,et al.FEBS Lett. 358:219-224(1995).


Images



Western blot analysis of GLP-1 Receptor expression in Hela (A), mouse brain (B), rat brain (C) whole cell lysates. (Predicted band size: 53 kDa; Observed band size: 53 kDa)



Immunohistochemical analysis of GLP-1 Receptor staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of GLP-1 Receptor staining in SKNSH cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.