

Anti-GRASP55 Antibody

Catalog # AP51234

Product Information

Application	WB, IHC
Primary Accession	Q9H8Y8
Reactivity	Human, Mouse, Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	47145

Additional Information

Gene ID	26003
Other Names	GOLPH6; Golgi reassembly-stacking protein 2; GRS2; Golgi phosphoprotein 6; GOLPH6; Golgi reassembly-stacking protein of 55 kDa; GRASP55; p59
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human GRASP55. The exact sequence is proprietary.
Dilution	WB~~ 1:1000 IHC~~1:100~500
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	GORASP2
Synonyms	GOLPH6
Function	Key structural protein of the Golgi apparatus (PubMed: 33301566). The membrane cisternae of the Golgi apparatus adhere to each other to form stacks, which are aligned side by side to form the Golgi ribbon (PubMed: 33301566). Acting in concert with GORASP1/GRASP65, is required for the formation and maintenance of the Golgi ribbon, and may be dispensable for the formation of stacks (PubMed: 33301566). However, other studies suggest that GORASP2 plays a role in the assembly and membrane stacking of the Golgi cisternae, and in the process by which Golgi stacks reform after breakdown during mitosis and meiosis (PubMed: 10487747 , PubMed: 21515684 , PubMed: 22523075). May regulate the intracellular transport and presentation of a defined set of transmembrane proteins, such as transmembrane TGFA (PubMed: 11101516). Required for normal acrosome formation during spermiogenesis and normal male fertility, probably by

promoting colocalization of JAM2 and JAM3 at contact sites between germ cells and Sertoli cells (By similarity). Mediates ER stress-induced unconventional (ER/Golgi-independent) trafficking of core-glycosylated CFTR to cell membrane (PubMed:[21884936](#), PubMed:[27062250](#), PubMed:[28067262](#)).


Cellular Location

Golgi apparatus membrane; Lipid-anchor. Endoplasmic reticulum membrane. Golgi apparatus. Note=Detected in the intermediate Golgi, membrane-associated (By similarity). ER stress triggers its relocation from Golgi to ER membrane (PubMed:27062250, PubMed:28067262). {ECO:0000250|UniProtKB:Q9R064, ECO:0000269|PubMed:27062250, ECO:0000269|PubMed:28067262}


References

Kuo A.,et al.EMBO J. 19:6427-6439(2000).
Jesch S.A.,et al.Mol. Biol. Cell 12:1811-1817(2001).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Hillier L.W.,et al.Nature 434:724-731(2005).

Images



Western blot analysis of GRASP55 expression in Hela (A), HGC27 (B), mouse testis (C), rat liver (D), rat testis (E) whole cell lysates. (Predicted band size: 47 kD; Observed band size: 55 kD)



Immunohistochemical analysis of GRASP55 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.