

GRK6 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51246

Product Information

| | |
|--------------------------|------------------------|
| Application | WB |
| Primary Accession | P43250 |
| Reactivity | Human, Mouse, Rat |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 65991 |

Additional Information

| | |
|---------------------------|--|
| Gene ID | 2870 |
| Other Names | G protein-coupled receptor kinase 6, G protein-coupled receptor kinase GRK6, GRK6, GPRK6 |
| Target/Specificity | KLH conjugated synthetic peptide derived from human GRK6 |
| Dilution | WB~~ 1:1000 |
| Format | 0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50% |
| Storage | Store at -20 °C.Stable for 12 months from date of receipt |

Protein Information

| | |
|--------------------------|---|
| Name | GRK6 |
| Synonyms | GPRK6 |
| Function | Specifically phosphorylates the activated forms of G protein- coupled receptors. Such receptor phosphorylation initiates beta- arrestin-mediated receptor desensitization, internalization, and signaling events leading to their desensitization. Seems to be involved in the desensitization of D2-like dopamine receptors in striatum and chemokine receptor CXCR4 which is critical for CXCL12-induced cell chemotaxis (By similarity). Phosphorylates rhodopsin (RHO) (in vitro) and a non G-protein-coupled receptor: LRP6 during Wnt signaling (in vitro). |
| Cellular Location | Membrane; Lipid-anchor. |
| Tissue Location | Widely expressed.. |


Background

Specifically phosphorylates the activated forms of G protein-coupled receptors. Such receptor phosphorylation initiates beta-arrestin-mediated receptor desensitization, internalization, and signaling events leading to their desensitization. Seems to be involved in the desensitization of D2-like dopamine receptors in striatum and chemokine receptor CXCR4 which is critical for CXCL12-induced cell chemotaxis (By similarity). Phosphorylates rhodopsin (RHO) (in vitro) and a non G-protein-coupled receptor: LRP6 during Wnt signaling (in vitro).

References

Benovic J.L.,et al.J. Biol. Chem. 268:19521-19527(1993).
Premont R.T.,et al.J. Biol. Chem. 274:29381-29389(1999).
Schmutz J.,et al.Nature 431:268-274(2004).
Haribabu B.,et al.Proc. Natl. Acad. Sci. U.S.A. 90:9398-9402(1993).
Stoffel R.H.,et al.J. Biol. Chem. 269:27791-27794(1994).

Images



All lanes : Anti-GRK6 Antibody at 1:1000 dilution Lane 1: HeLa whole cell lysates Lane 2: human brain lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 66 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.