

# mGLUR8 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51249

#### **Product Information**

Application WB Primary Accession 000222

**Reactivity** Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW101741

### **Additional Information**

**Gene ID** 2918

Other Names Metabotropic glutamate receptor 8, mGluR8, GRM8, GPRC1H, MGLUR8

**Dilution** WB~~1:1000

Format 0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

**Storage** Store at -20 °C.Stable for 12 months from date of receipt

## **Protein Information**

Name GRM8

**Synonyms** GPRC1H, MGLUR8

**Function** G-protein coupled receptor for glutamate. Ligand binding causes a

conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors, such as adenylate cyclase. Signaling inhibits adenylate cyclase activity.

**Cellular Location** Cell membrane; Multi-pass membrane protein.

# **Background**

G-protein coupled receptor for glutamate. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors, such as adenylate cyclase. Signaling inhibits adenylate cyclase activity.

#### References

Scherer S.W.,et al.Genomics 44:232-236(1997). Wu S.,et al.Brain Res. Mol. Brain Res. 53:88-97(1998). Malherbe P.,et al.Brain Res. Mol. Brain Res. 67:201-210(1999). Stormann T.M.,et al.Submitted (APR-2004) to the EMBL/GenBank/DDBJ databases. Kaighin V.A.,et al.Submitted (DEC-2007) to the EMBL/GenBank/DDBJ databases.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.