

# Histone H4 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51257

## Product Information

---

Application	WB
Primary Accession	<a href="#">P62805</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	11367

## Additional Information

---

Gene ID	121504;554313;8294;8359;8360;8361;8362;8363;8364;8365;8366;8367;8368;8370
Other Names	Histone H4, HIST1H4A, H4/A, H4FA
Target/Specificity	KLH conjugated synthetic peptide derived from human Histone H4
Dilution	WB~~ 1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

---

Name	H4C1
Synonyms	H4/A, H4FA, HIST1H4A
Function	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Cellular Location	Nucleus {ECO:0000250 UniProtKB:P62806}. Chromosome. Note=Localized to the nucleus when acetylated in step 11 spermatids. {ECO:0000250 UniProtKB:P62806}

## Background

---

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.


## References

---

Sierra F.,et al.Nucleic Acids Res. 11:7069-7086(1983).  
Pauli U.,et al.Science 236:1308-1311(1987).  
Albig W.,et al.Genomics 10:940-948(1991).  
Drabent B.,et al.DNA Cell Biol. 14:591-597(1995).  
Albig W.,et al.Gene 184:141-148(1997).

## Images

---



All lanes : Anti-Histone H4 Antibody at 1:1000 dilution  
Lane 1: HeLa whole cell lysates Lane 2: HepG2 whole cell  
lysates Lysates/proteins at 20 µg per lane. Secondary  
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at  
1/10000 dilution Predicted band size : 11 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.