

KAT7 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51295

Product Information

Application	WB
Primary Accession	O95251
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	70642

Additional Information

Gene ID	11143
Other Names	Histone acetyltransferase KAT7, Histone acetyltransferase binding to ORC1, Lysine acetyltransferase 7, MOZ, YBF2/SAS3, SAS2 and TIP60 protein 2, MYST-2, KAT7, HBO1, HBOa, MYST2
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human KAT7. The exact sequence is proprietary.
Dilution	WB~~ 1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	KAT7 {ECO:0000303 PubMed:31767635, ECO:0000312 HGNC:HGNC:17016}
Function	Catalytic subunit of histone acetyltransferase HBO1 complexes, which specifically mediate acetylation of histone H3 at 'Lys-14' (H3K14ac), thereby regulating various processes, such as gene transcription, protein ubiquitination, immune regulation, stem cell pluripotent and self-renewal maintenance and embryonic development (PubMed: 16387653 , PubMed: 21753189 , PubMed: 24065767 , PubMed: 26620551 , PubMed: 31767635 , PubMed: 31827282). Some complexes also catalyze acetylation of histone H4 at 'Lys-5', 'Lys-8' and 'Lys-12' (H4K5ac, H4K8ac and H4K12ac, respectively), regulating DNA replication initiation, regulating DNA replication initiation (PubMed: 10438470 , PubMed: 19187766 , PubMed: 20129055 , PubMed: 24065767). Specificity of the HBO1 complexes is determined by the scaffold subunit: complexes containing BRPF scaffold (BRPF1, BRD1/BRPF2 or BRPF3) direct KAT7/HBO1 specificity towards H3K14ac, while complexes containing JADE (JADE1, JADE2 and JADE3) scaffold

direct KAT7/HBO1 specificity towards histone H4 (PubMed:[19187766](#), PubMed:[20129055](#), PubMed:[24065767](#), PubMed:[26620551](#)). H3K14ac promotes transcriptional elongation by facilitating the processivity of RNA polymerase II (PubMed:[31827282](#)). Acts as a key regulator of hematopoiesis by forming a complex with BRD1/BRPF2, directing KAT7/HBO1 specificity towards H3K14ac and promoting erythroid differentiation (PubMed:[21753189](#)). H3K14ac is also required for T-cell development (By similarity). KAT7/HBO1-mediated acetylation facilitates two consecutive steps, licensing and activation, in DNA replication initiation: H3K14ac facilitates the activation of replication origins, and histone H4 acetylation (H4K5ac, H4K8ac and H4K12ac) facilitates chromatin loading of MCM complexes, promoting DNA replication licensing (PubMed:[10438470](#), PubMed:[11278932](#), PubMed:[18832067](#), PubMed:[19187766](#), PubMed:[20129055](#), PubMed:[21856198](#), PubMed:[24065767](#), PubMed:[26620551](#)). Acts as a positive regulator of centromeric CENPA assembly: recruited to centromeres and mediates histone acetylation, thereby preventing centromere inactivation mediated by SUV39H1, possibly by increasing histone turnover/exchange (PubMed:[27270040](#)). Involved in nucleotide excision repair: phosphorylation by ATR in response to ultraviolet irradiation promotes its localization to DNA damage sites, where it mediates histone acetylation to facilitate recruitment of XPC at the damaged DNA sites (PubMed:[28719581](#)). Acts as an inhibitor of NF-kappa-B independently of its histone acetyltransferase activity (PubMed:[16997280](#)).

Cellular Location

Nucleus. Chromosome. Chromosome, centromere. Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q5SVQ0}. Note=Associates with replication origins specifically during the G1 phase of the cell cycle (PubMed:18832067, PubMed:20129055). Localizes to transcription start sites (PubMed:21753189, PubMed:24065767). Localizes to ultraviolet- induced DNA damage sites following phosphorylation by ATR (PubMed:28719581). Localizes to centromeres in G1 phase (PubMed:27270040).

Tissue Location

Ubiquitously expressed, with highest levels in testis.

Background


Component of the HBO1 complex which has a histone H4- specific acetyltransferase activity, a reduced activity toward histone H3 and is responsible for the bulk of histone H4 acetylation in vivo. Through chromatin acetylation it may regulate DNA replication and act as a coactivator of TP53-dependent transcription. Specifically represses AR-mediated transcription.

References

Iizuka M.,et al.J. Biol. Chem. 274:23027-23034(1999).
 Sharma M.,et al.J. Biol. Chem. 275:35200-35208(2000).
 Jian J.,et al.Submitted (APR-1999) to the EMBL/GenBank/DDBJ databases.
 Borrow J.,et al.Submitted (DEC-1999) to the EMBL/GenBank/DDBJ databases.
 Ota T.,et al.Nat. Genet. 36:40-45(2004).

Images

All lanes : Anti-KAT7 Antibody at 1:1000 dilution Lane 1:
 Jurkat whole cell lysates Lane 2: Hela whole cell lysates
 Lysates/proteins at 20 µg per lane. Secondary Goat
 Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000



dilution Predicted band size : 71 kDa Blocking/Dilution
buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.