10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Cytokeratin 10 Antibody

Catalog # AP51305

Product Information

Application	WB, IHC, IF/IC
Primary Accession	P13645
Reactivity	Human, Mouse, Rat, Rabbit, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	58827

Additional Information

Gene ID	3858
Other Names	KPP; Keratin, type I cytoskeletal 10; Cytokeratin-10; CK-10; Keratin-10; K10
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Cytokeratin 10. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	KRT10 (HGNC:6413)
Synonyms	KPP
Function	Structural component of intermediate filaments in suprabasal keratinocytes of stratified epithelia. Forms heteropolymers primarily with the type II keratin KRT1, and also with KRT2, assembling into intermediate filament networks that support terminal differentiation, mechanical resilience, and epidermal barrier formation.
Cellular Location	Secreted, extracellular space. Cell surface. Cytoplasm
Tissue Location	Seen in all suprabasal cell layers including stratum corneum. Expressed on the surface of lung cell lines (PubMed:19627498). Localized on the surface of desquamated nasal epithelial cells (at protein level) (PubMed:12427098)


References

Zhou X.M.,et al.J. Biol. Chem. 263:15584-15589(1988).
Rieger M.,et al.J. Mol. Biol. 204:841-856(1988).
Zody M.C.,et al.Nature 440:1045-1049(2006).
Darmon M.Y.,et al.Mol. Biol. Rep. 12:277-283(1987).
Rothnagel J.A.,et al.J. Invest. Dermatol. 102:13-16(1994).


Images



Western blot analysis of Cytokeratin 10 expression in HeLa (A), A431 (B), Raw264.7 (C), NIHC3T3 (D), rat brain (E) whole cell lysates. (Predicted band size: 58 kD; Observed band size: 60 kD)



Immunohistochemical analysis of Cytokeratin 10 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of Cytokeratin 10 staining in A431 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.