10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Cytokeratin 14 Antibody

Catalog # AP51307

Product Information

Application	WB, IHC, IF/IC
Primary Accession	P02533
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51561

Additional Information

Gene ID	3861
Other Names	KRT14; Keratin, type I cytoskeletal 14; Cytokeratin-14; CK-14; Keratin-14; K14
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human Cytokeratin 14. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	KRT14 (HGNC:6416)
Function	Structural component of intermediate filaments in basal keratinocytes of stratified epithelia. Forms heteropolymers with type II keratin KRT5, contributing to the keratin intermediate filament network that confers mechanical strength and structural integrity to the basal layer of the epidermis and other stratified epithelial (PubMed:11724817, PubMed:1694855, PubMed:1720261, PubMed:41511042). The nonhelical tail domain is involved in promoting KRT5-KRT14 filaments to self-organize into large bundles and enhances the mechanical properties involved in resilience of keratin intermediate filaments in vitro (PubMed:11724817). Displays an antagonist role with KRT15 in basal epithelial cells, where KRT14 promotes cell differentiation while KRT15 maintains the progenitor state and promotes proliferation and clonogenicity (PubMed:41511042). Promotes cell differentiation by retaining YAP1 in the cytoplasm of basal keratinocytes via regulatory residue C367 and stratifin/SNF interaction (PubMed:41511042). Contributes to airway basal cell differentiation after injury (PubMed:36512409).

Cellular Location	Cytoplasm. Nucleus. Note=Expressed in both as a filamentous pattern.
Tissue Location	Expressed in the corneal epithelium (at protein level) (PubMed:26758872). Detected in the basal layer, lowered within the more apically located layers specifically in the stratum spinosum, stratum granulosum but is not detected in stratum corneum. Strongly expressed in the outer root sheath of anagen follicles but not in the germinative matrix, inner root sheath or hair (PubMed:9457912). Found in keratinocytes surrounding the club hair during telogen (PubMed:9457912). Expressed in gland ducts and in the myoepithelial cells of gland tubules and acini in pulmonary bronchus (PubMed:36512409).


References

Marchuk D.,et al.Cell 39:491-498(1984).
Marchuk D.,et al.Proc. Natl. Acad. Sci. U.S.A. 82:1609-1613(1985).
Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
Zody M.C.,et al.Nature 440:1045-1049(2006).
Hanukoglu I.,et al.Cell 31:243-252(1982).

Images




Western blot analysis of Cytokeratin 14 expression in MCF7 (A), mouse brain (B), rat brain (C) whole cell lysates. (Predicted band size: 51 kD; Observed band size: 52 kD)



Immunohistochemical analysis of Cytokeratin 14 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Immunofluorescent analysis of Cytokeratin 14 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.