10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-EDG7 Antibody

Catalog # AP51321

Product Information

Application	WB, IHC
Primary Accession	Q9UBY5
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40128

Additional Information

Gene ID	23566
Other Names	EDG7; LPA3; Lysophosphatidic acid receptor 3; LPA receptor 3; LPA-3; Lysophosphatidic acid receptor Edg-7
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human EDG7. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	LPAR3
Synonyms	EDG7, LPA3
Function	Receptor for lysophosphatidic acid (LPA), a mediator of diverse cellular activities. May play a role in the development of ovarian cancer. Seems to be coupled to the G(i)/G(o) and G(q) families of heteromeric G proteins.
Cellular Location	Cell membrane; Multi-pass membrane protein.
Tissue Location	Most abundantly expressed in prostate, testes, pancreas, and heart, with moderate levels in lung and ovary. No detectable expression in brain, placenta, liver, skeletal muscle, kidney, spleen, thymus, small intestine, colon, or peripheral blood leukocytes


References

Bandoh K.,et al.J. Biol. Chem. 274:27776-27785(1999).
Im D.-S.,et al.Mol. Pharmacol. 57:753-759(2000).
Kopatz S.A.,et al.Submitted (JUN-2003) to the EMBL/GenBank/DDBJ databases.
Contos J.J.A.,et al.Mol. Pharmacol. 58:1188-1196(2000).

Images



Western blot analysis of EDG7 expression in HepG2 (A), NT2D1 (B) whole cell lysates. (Predicted band size: 40 kD; Observed band size: 40 kD)



Immunohistochemical analysis of EDG7 staining in human liver cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.