10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Anti-JIP2 Antibody

Catalog # AP51328

## Product Information

| | |
|---|---|
| **Application** | WB, IHC, IF/IC |
| **Primary Accession** | Q13387 |
| **Reactivity** | Human, Mouse, Rat, Zebrafish |
| **Host** | Rabbit |
| **Clonality** | Polyclonal |
| **Calculated MW** | 87975 |

## Additional Information


| | |
|---|---|
| **Gene ID** | 23542 |
| **Other Names** | IB2; JIP2; PRKM8IPL; C-Jun-amino-terminal kinase-interacting protein 2; JIP-2; JNK-interacting protein 2; Islet-brain-2; IB-2; JNK MAP kinase scaffold protein 2; Mitogen-activated protein kinase 8-interacting protein 2 |
| **Target/Specificity** | KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human JIP2. The exact sequence is proprietary. |
| **Dilution** | WB~~WB (1/500 - 1/1000) IHC~~1:100~500 IF/IC~~N/A |
| **Format** | Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide. |
| **Storage** | Store at -20 °C.Stable for 12 months from date of receipt |

## Protein Information


| | |
|---|---|
| **Name** | MAPK8IP2 |
| **Synonyms** | IB2, JIP2, PRKM8IPL |
| **Function** | The JNK-interacting protein (JIP) group of scaffold proteins selectively mediates JNK signaling by aggregating specific components of the MAPK cascade to form a functional JNK signaling module. JIP2 inhibits IL1 beta-induced apoptosis in insulin-secreting cells. May function as a regulator of vesicle transport, through interactions with the JNK-signaling components and motor proteins (By similarity). |
| **Cellular Location** | Cytoplasm. Note=Accumulates in cell surface projections |
| **Tissue Location** | Expressed mainly in the brain and pancreas, including insulin-secreting cells. In the nervous system, more abundantly expressed in the cerebellum, pituitary gland, occipital lobe and the amygdala. Also expressed in fetal brain. |

Very low levels found in uterus, ovary, prostate, colon, testis, adrenal gland, thyroid gland and salivary gland


## Images




Western blot analysis of JIP2 expression in SHSY5Y (A), H1688 (B), mouse liver (C), rat liver (D), mouse testis (E), rat testis (F) whole cell lysates. (Predicted band size: 87 kD; Observed band size: 100 kD)



Immunohistochemical analysis of JIP2 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of JIP2 staining in Raw264.7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.