

MMP14 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51349

Product Information

Application	WB
Primary Accession	P50281
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	65894

Additional Information

Gene ID	4323
Other Names	Matrix metalloproteinase-14, MMP-14, MMP-X1, Membrane-type matrix metalloproteinase 1, MT-MMP 1, MTMMP1, Membrane-type-1 matrix metalloproteinase, MT1-MMP, MT1MMP, MMP14
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human MMP14. The exact sequence is proprietary.
Dilution	WB~~ 1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	MMP14
Function	Endopeptidase that degrades various components of the extracellular matrix such as collagen (PubMed: 8015608). Essential for pericellular collagenolysis and modeling of skeletal and extraskelatal connective tissues during development (By similarity). Activates progelatinase A/MMP2, thereby acting as a positive regulator of cell growth and migration (PubMed: 22065321 , PubMed: 8015608). Involved in the formation of the fibrovascular tissues in association with pro-MMP2 (PubMed: 12714657 , PubMed: 22065321). May be involved in actin cytoskeleton reorganization by cleaving PTK7 (PubMed: 20837484). Acts as a regulator of Notch signaling by mediating cleavage and inhibition of DLL1 (PubMed: 21572390). Cleaves ADGRB1 to release vasculostatin-40 which inhibits angiogenesis (PubMed: 22330140). Acts as a negative regulator of the GDF15-GFRAL aversive response by mediating cleavage and inactivation of GFRAL (PubMed: 35177851).

Cellular Location	Cell membrane; Single-pass type I membrane protein. Melanosome. Cytoplasm Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065). Forms a complex with BST2 and localizes to the cytoplasm (PubMed:17081065)
Tissue Location	Expressed in stromal cells of colon, breast, and head and neck. Expressed in lung tumors.


Background

Seems to specifically activate progelatinase A. May thus trigger invasion by tumor cells by activating progelatinase A on the tumor cell surface. May be involved in actin cytoskeleton reorganization by cleaving PTK7. Acts as a positive regulator of cell growth and migration via activation of MMP15.

References

Sato H.,et al.Nature 370:61-65(1994).
Takino T.,et al.Gene 155:293-298(1995).
Okada A.,et al.Proc. Natl. Acad. Sci. U.S.A. 92:2730-2734(1995).
Will H.,et al.Eur. J. Biochem. 231:602-608(1995).
Luo G.-X.,et al.Submitted (NOV-1995) to the EMBL/GenBank/DDBJ databases.

Images



Anti-MMP14 Antibody at 1:1000 dilution + human breast lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 66 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.