

Anti-NEIL3 Antibody

Catalog # AP51383

Product Information

Application	WB, IHC
Primary Accession	Q8TAT5
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67769

Additional Information

Gene ID	55247
Other Names	Endonuclease 8-like 3; DNA glycosylase FPG2; DNA glycosylase/AP lyase Neil3; Endonuclease VIII-like 3; Nei-like protein 3
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human NEIL3. The exact sequence is proprietary.
Dilution	WB~~ 1:1000 IHC~~1:100~500
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	NEIL3
Function	DNA glycosylase which prefers single-stranded DNA (ssDNA), or partially ssDNA structures such as bubble and fork structures, to double-stranded DNA (dsDNA) (PubMed: 12433996 , PubMed: 19170771 , PubMed: 22569481 , PubMed: 23755964). Mediates interstrand cross-link repair in response to replication stress: acts by mediating DNA glycosylase activity, cleaving one of the two N-glycosyl bonds comprising the interstrand cross-link, which avoids the formation of a double-strand break but generates an abasic site that is bypassed by translesion synthesis polymerases (By similarity). In vitro, displays strong glycosylase activity towards the hydantoin lesions spiroiminodihydantoin (Sp) and guanidinohydantoin (Gh) in both ssDNA and dsDNA; also recognizes FapyA, FapyG, 5-OHU, 5-OHC, 5-OHMH, Tg and 8-oxoA lesions in ssDNA (PubMed: 12433996 , PubMed: 19170771 , PubMed: 22569481 , PubMed: 23755964). No activity on 8-oxoG detected (PubMed: 12433996 , PubMed: 19170771 , PubMed: 22569481 , PubMed: 23755964). Also shows weak DNA-(apurinic or apyrimidinic site) lyase activity (PubMed: 12433996 ,

PubMed:[19170771](#), PubMed:[22569481](#), PubMed:[23755964](#)). In vivo, appears to be the primary enzyme involved in removing Sp and Gh from ssDNA in neonatal tissues (PubMed:[12433996](#), PubMed:[19170771](#), PubMed:[22569481](#), PubMed:[23755964](#)).

Cellular Location

Nucleus. Chromosome {ECO:0000250|UniProtKB:A0A1L8HU22}.
Note=Recruited to replication stress sites via interaction with ubiquitinated CMG helicase {ECO:0000250|UniProtKB:A0A1L8HU22}


Tissue Location

Expressed in keratinocytes and embryonic fibroblasts (at protein level). Also detected in thymus, testis and fetal lung primary fibroblasts.


References

Takao M.,et al.J. Biol. Chem. 277:42205-42213(2002).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Hillier L.W.,et al.Nature 434:724-731(2005).
Morland I.,et al.Nucleic Acids Res. 30:4926-4936(2002).
Torisu K.,et al.J. Biochem. 138:763-772(2005).

Images



Western blot analysis of NEIL3 expression in MCF7 (A), H1792 (B), rat thymus (C), mouse testis (D) whole cell lysates. (Predicted band size: 67 kD; Observed band size: 68 kD)



Immunohistochemical analysis of NEIL3 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.