

# NOTCH2 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51394

## **Product Information**

Application	WB, ICC, IHC-P
Primary Accession	<u>Q04721</u>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	265405

## **Additional Information**

Gene ID	4853
Other Names	Neurogenic locus notch homolog protein 2, Notch 2, hN2, Notch 2 extracellular truncation, Notch 2 intracellular domain, NOTCH2
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human NOTCH2. The exact sequence is proprietary.
Dilution	WB~~1:1000 ICC~~N/A IHC-P~~N/A
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

#### **Protein Information**

Name	NOTCH2 ( <u>HGNC:7882</u> )
Function	Functions as a receptor for membrane-bound ligands Jagged-1 (JAG1), Jagged-2 (JAG2) and Delta-1 (DLL1) to regulate cell-fate determination. Upon ligand activation through the released notch intracellular domain (NICD) it forms a transcriptional activator complex with RBPJ/RBPSUH and activates genes of the enhancer of split locus (PubMed: <u>21378985</u> , PubMed: <u>21378989</u> ). Affects the implementation of differentiation, proliferation and apoptotic programs (By similarity). Involved in bone remodeling and homeostasis. In collaboration with RELA/p65 enhances NFATc1 promoter activity and positively regulates RANKL-induced osteoclast differentiation (PubMed: <u>29149593</u> ). Positively regulates self-renewal of liver cancer cells (PubMed: <u>25985737</u> ).
Cellular Location	[Notch 2 extracellular truncation]: Cell membrane; Single-pass type I membrane protein

## Background

Functions as a receptor for membrane-bound ligands Jagged1, Jagged2 and Delta1 to regulate cell-fate determination. Upon ligand activation through the released notch intracellular domain (NICD) it forms a transcriptional activator complex with RBPJ/RBPSUH and activates genes of the enhancer of split locus. Affects the implementation of differentiation, proliferation and apoptotic programs (By similarity). Involved in bone remodeling and homeostasis. In collaboration with RELA/p65 enhances NFATc1 promoter activity and positively regulates RANKL-induced osteoclast differentiation.

#### References

Blaumueller C.M.,et al.Submitted (NOV-2000) to the EMBL/GenBank/DDBJ databases. Correa R.G.,et al.Submitted (OCT-2000) to the EMBL/GenBank/DDBJ databases. Gregory S.G.,et al.Nature 441:315-321(2006). Lemasson I.,et al.Submitted (NOV-1996) to the EMBL/GenBank/DDBJ databases. Stifani S.,et al.Nat. Genet. 2:119-127(1992).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.