10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-NOTCH2 Antibody

Catalog # AP51394

Product Information

Application	WB, IHC
Primary Accession	Q04721
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	265405

Additional Information

Gene ID	4853
Other Names	Neurogenic locus notch homolog protein 2; Notch 2; hN2
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human NOTCH2. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	NOTCH2 (HGNC:7882)
Function	Functions as a receptor for membrane-bound ligands Jagged-1 (JAG1), Jagged-2 (JAG2) and Delta-1 (DLL1) to regulate cell-fate determination. Upon ligand activation through the released notch intracellular domain (NICD) it forms a transcriptional activator complex with RBPJ/RBPSUH and activates genes of the enhancer of split locus (PubMed:21378985, PubMed:21378989). Affects the implementation of differentiation, proliferation and apoptotic programs (By similarity). Involved in bone remodeling and homeostasis. In collaboration with RELA/p65 enhances NFATc1 promoter activity and positively regulates RANKL-induced osteoclast differentiation (PubMed:29149593). Positively regulates self-renewal of liver cancer cells (PubMed:25985737).
Cellular Location	[Notch 2 extracellular truncation]: Cell membrane; Single-pass type I membrane protein
Tissue Location	Expressed in the brain, heart, kidney, lung, skeletal muscle and liver.

Ubiquitously expressed in the embryo


References

Blaumueller C.M.,et al.Submitted (NOV-2000) to the EMBL/GenBank/DDBJ databases.
Correa R.G.,et al.Submitted (OCT-2000) to the EMBL/GenBank/DDBJ databases.
Gregory S.G.,et al.Nature 441:315-321(2006).
Lemasson I.,et al.Submitted (NOV-1996) to the EMBL/GenBank/DDBJ databases.
Stifani S.,et al.Nat. Genet. 2:119-127(1992).

Images



Western blot analysis of NOTCH2 expression in HEK293T (A), HepG2 (B) whole cell lysates. (Predicted band size: 265 kD; Observed band size: 110 kD)



Immunohistochemical analysis of NOTCH2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.