

OTUB2 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51410

Product Information

Application	WB
Primary Accession	Q96DC9
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	27213

Additional Information

Gene ID	78990
Other Names	Ubiquitin thioesterase OTUB2, Deubiquitinating enzyme OTUB2, OTU domain-containing ubiquitin aldehyde-binding protein 2, Otubain-2, Ubiquitin-specific-processing protease OTUB2, OTUB2, C14orf137, OTB2, OTU2
Target/Specificity	KLH conjugated synthetic peptide derived from human OTUB2
Dilution	WB~~ 1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	OTUB2
Synonyms	C14orf137, OTB2, OTU2
Function	Hydrolase that can remove conjugated ubiquitin from proteins in vitro and may therefore play an important regulatory role at the level of protein turnover by preventing degradation. Mediates deubiquitination of 'Lys-11'-, 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains, with a preference for 'Lys-63'-linked polyubiquitin chains.
Tissue Location	Widely expressed. Expressed at higher level in brain.


Background

Hydrolase that can remove conjugated ubiquitin from proteins in vitro and may therefore play an important regulatory role at the level of protein turnover by preventing degradation. Mediates deubiquitination of 'Lys-11'-, 'Lys-48'- and 'Lys-63'- linked polyubiquitin chains, with a preference for 'Lys-63'-linked polyubiquitin chains.

References

Balakirev M.Y., et al. EMBO Rep. 4:517-522(2003).
Ota T., et al. Nat. Genet. 36:40-45(2004).
Ebert L., et al. Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.
Heilig R., et al. Nature 421:601-607(2003).
Edelmann M.J., et al. Biochem. J. 418:379-390(2009).

Images



All lanes : Anti-OTUB2 Antibody at 1:1000 dilution Lane 1: HT-29 whole cell lysates Lane 2: PC-3 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 27 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.