

Anti-OCT2 Antibody

Catalog # AP51440

Product Information

Application	WB, IHC, IF/IC
Primary Accession	P09086
Reactivity	Human, Mouse, Rat, Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51209

Additional Information

Gene ID	5452
Other Names	OCT2; OTF2; POU domain, class 2, transcription factor 2; Lymphoid-restricted immunoglobulin octamer-binding protein NF-A2; Octamer-binding protein 2; Oct-2; Octamer-binding transcription factor 2; OTF-2
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human OCT2. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	POU2F2 (HGNC:9213)
Synonyms	OCT2, OTF2
Function	Transcription factor that specifically binds to the octamer motif (5'-ATTGTCAT-3') (PubMed: 2904654 , PubMed: 7859290). Regulates IL6 expression in B cells with POU2AF1 (By similarity). Regulates transcription in a number of tissues in addition to activating immunoglobulin gene expression (PubMed: 2901913 , PubMed: 2904654). Modulates transcription transactivation by NR3C1, AR and PGR (PubMed: 10480874).
Cellular Location	Cytoplasm. Nucleus.
Tissue Location	Isoform 3 is B-cell specific. Isoform 5 is expressed in B-cells and the immunoglobulin-expressing T-cell line MOLT-4, but not in the T-cell line BW5147.


References

Clerc R.G.,et al.Genes Dev. 2:1570-1581(1988).
Scheidereit C.,et al.Nature 336:551-557(1988).
Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
Mueller M.M.,et al.Nature 336:544-551(1988).
Muller-Immergluck M.M.,et al.EMBO J. 9:1625-1634(1990).


Images



Western blot analysis of OCT2 expression in rat spleen (A), mouse spleen (B), K562 (C), A2780 (D), Myla2059 (E) whole cell lysates. (Predicted band size: 51 kDa; Observed band size: 75 kDa)



Immunohistochemical analysis of OCT2 staining in human tonsil formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of OCT2 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.