

Anti-Beta-glucosidase 2 Antibody

Catalog # AP51448

Product Information

Application	WB, IHC, IF/IC
Primary Accession	P14314
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59425

Additional Information

Gene ID	5589
Other Names	G19P1; Glucosidase 2 subunit beta; 80K-H protein; Glucosidase II subunit beta; Protein kinase C substrate 60.1 kDa protein heavy chain; PKCSH
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Beta-glucosidase 2. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	PRKCSH {ECO:0000303 PubMed:28375157, ECO:0000312 HGNC:HGNC:9411}
Function	Regulatory subunit of glucosidase II that cleaves sequentially the 2 innermost alpha-1,3-linked glucose residues from the Glc(2)Man(9)GlcNAc(2) oligosaccharide precursor of immature glycoproteins (PubMed: 10929008). Required for efficient PKD1/Polycystin-1 biogenesis and trafficking to the plasma membrane of the primary cilia (By similarity).
Cellular Location	Endoplasmic reticulum {ECO:0000255 PROSITE- ProRule:PRU10138, ECO:0000305 PubMed:10929008}

References


Sakai K.,et al.Genomics 5:309-315(1989).

Ophoff R.A., et al. *Eur. J. Hum. Genet.* 4:321-328(1996).
 Pelletier M.F., et al. *Glycobiology* 10:815-827(2000).
 Kalnine N., et al. Submitted (AUG-2003) to the EMBL/GenBank/DDBJ databases.
 Ota T., et al. *Nat. Genet.* 36:40-45(2004).


Images



Western blot analysis of Beta-glucosidase 2 expression in A375 (A), MCF7 (B), mouse brain (C) whole cell lysates. (Predicted band size: 59 kD; Observed band size: 87 kD)



Immunohistochemical analysis of Beta-glucosidase 2 staining in human colorectal cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of Beta-glucosidase 2 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.