

Anti-RPL17 Antibody

Catalog # AP51482

Product Information

Application	WB, IHC, IF/IC
Primary Accession	P18621
Reactivity	Human, Mouse, Rat, Bovine, Mk
Host	Rabbit
Clonality	Polyclonal
Calculated MW	21397

Additional Information

Gene ID	6139
Other Names	60S ribosomal protein L17; 60S ribosomal protein L23; PD-1
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human RPL17. The exact sequence is proprietary.
Dilution	WB~~ 1:1000 IHC~~1:100~500 IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	RPL17
Function	Component of the large ribosomal subunit (PubMed: 12962325 , PubMed: 23636399 , PubMed: 32669547). The ribosome is a large ribonucleoprotein complex responsible for the synthesis of proteins in the cell (PubMed: 12962325 , PubMed: 23636399 , PubMed: 32669547).
Cellular Location	Cytoplasm.
Tissue Location	Expressed in pancreas, lung, colon, cystic duct, gall bladder, kidney and liver. Expressed at high levels in the well differentiated pancreatic tumor cell lines HPAF, COLO 357 and Capan-1, the moderately differentiated pancreatic tumor cell lines T3M-4, AsPc-1 and BxPc-3, the poorly differentiated pancreatic tumor cell line MIA PaCa-2, and the pancreatic tumor cell lines of undefined differentiation status such as SW979. Expressed at lower levels in the poorly differentiated pancreatic tumor cell lines HCG-25 and PANC-1


References

Mager D.L.,et al.Nucleic Acids Res. 18:5301-5301(1990).
Batra S.K.,et al.Cell Growth Differ. 2:385-390(1991).
Yoshihama M.,et al.Genome Res. 12:379-390(2002).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Li W.B.,et al.Submitted (MAR-2004) to the EMBL/GenBank/DDBJ databases.


Images



Western blot analysis of RPL17 expression in A375 (A), Raw264.7 (B), H9C2 (C) whole cell lysates. (Predicted band size: 21 kD; Observed band size: 22 kD)



Immunohistochemical analysis of RPL17 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of RPL17 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.