

SAP155 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51507

Product Information

Application	WB
Primary Accession	<u>075533</u>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	145830

Additional Information

Gene ID	23451
Other Names	Splicing factor 3B subunit 1, Pre-mRNA-splicing factor SF3b 155 kDa subunit, SF3b155, Spliceosome-associated protein 155, SAP 155, SF3B1, SAP155
Target/Specificity	KLH conjugated synthetic peptide derived from human SAP155
Dilution	WB~~ 1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	SF3B1 {ECO:0000303 PubMed:30567737, ECO:0000312 HGNC:HGNC:10768}
Function	Component of the 17S U2 SnRNP complex of the spliceosome, a large ribonucleoprotein complex that removes introns from transcribed pre-mRNAs (PubMed:12234937, PubMed:27720643, PubMed:32494006, PubMed:34822310). The 17S U2 SnRNP complex (1) directly participates in early spliceosome assembly and (2) mediates recognition of the intron branch site during pre-mRNA splicing by promoting the selection of the pre-mRNA branch-site adenosine, the nucleophile for the first step of splicing (PubMed:32494006, PubMed:34822310). Within the 17S U2 SnRNP complex, SF3B1 is part of the SF3B subcomplex, which is required for 'A' complex assembly formed by the stable binding of U2 snRNP to the branchpoint sequence in pre-mRNA (PubMed:12234937). Sequence independent binding of SF3A and SF3B subcomplexes upstream of the branch site is essential, it may anchor U2 snRNP to the pre-mRNA (PubMed:12234937). May also be involved in the assembly of the 'E' complex (PubMed:10882114). Also acts as a component of the minor spliceosome, which is involved in the splicing of U12-type introns in pre-mRNAs (PubMed:15146077, PubMed:33509932).

	Together with other U2 snRNP complex components may also play a role in the selective processing of microRNAs (miRNAs) from the long primary miRNA transcript, pri-miR-17-92 (By similarity).
Cellular Location	Nucleus. Nucleus speckle. Note=During mitosis, transiently dispersed from the nuclear speckles to the cytoplasm

Background

Subunit of the splicing factor SF3B required for 'A' complex assembly formed by the stable binding of U2 snRNP to the branchpoint sequence (BPS) in pre-mRNA. Sequence independent binding of SF3A/SF3B complex upstream of the branch site is essential, it may anchor U2 snRNP to the pre-mRNA. May also be involved in the assembly of the 'E' complex. Belongs also to the minor U12-dependent spliceosome, which is involved in the splicing of rare class of nuclear pre-mRNA intron.

References

Wang C.,et al.Genes Dev. 12:1409-1414(1998). Woessner J.,et al.Submitted (SEP-1998) to the EMBL/GenBank/DDBJ databases. Hillier L.W.,et al.Nature 434:724-731(2005). Yu W.,et al.Submitted (JUN-1998) to the EMBL/GenBank/DDBJ databases. Das R.,et al.Mol. Cell 5:779-787(2000).

Images



All lanes : Anti-SAP155 Antibody at 1:1000 dilution Lane 1: Hela whole cell lysates Lane 2: Ramos whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L),Peroxidase conjugated at 1/10000 dilution Predicted band size : 146 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.