

Alpha-sarcoglycan Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51510

Product Information

Application WB Primary Accession Q16586

Reactivity Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW42875

Additional Information

Gene ID 6442

Other Names Alpha-sarcoglycan, Alpha-SG, 50 kDa dystrophin-associated glycoprotein,

50DAG, Adhalin, Dystroglycan-2, SGCA, ADL, DAG2

Target/Specificity KLH conjugated synthetic peptide derived from human Alpha-sarcoglycan

Dilution WB~~ 1:1000

Format 0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name SGCA

Synonyms ADL, DAG2

Function Component of the sarcoglycan complex, a subcomplex of the

dystrophin-glycoprotein complex which forms a link between the F-actin

cytoskeleton and the extracellular matrix.

Cellular Location Cell membrane, sarcolemma; Single- pass type I membrane protein.

Cytoplasm, cytoskeleton

Tissue Location Most strongly expressed in skeletal muscle. Also expressed in cardiac muscle

and, at much lower levels, in lung. In the fetus, most abundant in cardiac muscle and, at lower levels, in lung Also detected in liver and kidney. Not

expressed in brain

Background

Component of the sarcoglycan complex, a subcomplex of the dystrophin-glycoprotein complex which forms a link between the F-actin cytoskeleton and the extracellular matrix.

References

Roberds S.L.,et al.Cell 78:625-633(1994).


McNally E.,et al.Proc. Natl. Acad. Sci. U.S.A. 91:9690-9694(1994).

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Zody M.C.,et al.Nature 440:1045-1049(2006).

Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

Images



All lanes: Anti-Alpha-sarcoglycan Antibody at 1:1000 dilution Lane 1: human skeletal musle lysates Lane 2: human heart lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size: 43 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.