

ENT1 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51518

Product Information

Application	WB
Primary Accession	Q99808
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	50219

Additional Information

Gene ID	2030
Other Names	Equilibrative nucleoside transporter 1, Equilibrative nitrobenzylmercaptapurine riboside-sensitive nucleoside transporter, Equilibrative NBMPR-sensitive nucleoside transporter, Nucleoside transporter, es-type, Solute carrier family 29 member 1, SLC29A1, ENT1
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human ENT1. The exact sequence is proprietary.
Dilution	WB~~ 1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	SLC29A1 (HGNC:11003)
Synonyms	ENT1
Function	Uniporter involved in the facilitative transport of nucleosides and nucleobases, and contributes to maintaining their cellular homeostasis (PubMed: 10722669 , PubMed: 10755314 , PubMed: 12527552 , PubMed: 14759222 , PubMed: 15037197 , PubMed: 17379602 , PubMed: 21795683 , PubMed: 26406980 , PubMed: 27995448 , PubMed: 35790189 , PubMed: 8986748). Functions as a Na(+)-independent transporter (PubMed: 8986748). Involved in the transport of nucleosides such as adenosine, guanosine, inosine, uridine, thymidine and cytidine (PubMed: 10722669 , PubMed: 10755314 , PubMed: 12527552 , PubMed: 14759222 , PubMed: 15037197 , PubMed: 17379602 , PubMed: 26406980 , PubMed: 8986748). Also transports purine nucleobases

(hypoxanthine, adenine, guanine) and pyrimidine nucleobases (thymine, uracil) (PubMed:[21795683](#), PubMed:[27995448](#)). Mediates basolateral nucleoside uptake into Sertoli cells, thereby regulating the transport of nucleosides in testis across the blood-testis barrier (By similarity). Regulates inosine levels in brown adipocytes tissues (BAT) and extracellular inosine levels, which controls BAT-dependent energy expenditure (PubMed:[35790189](#)).

Cellular Location

Basolateral cell membrane; Multi-pass membrane protein. Apical cell membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein. Note=Localized to the basolateral membrane of Sertoli cells (PubMed:23639800). Localized to the cell membrane of erythrocytes (PubMed:11584005, PubMed:23219802).

Tissue Location

Expressed in testis at the blood-testis barrier (at protein level) (PubMed:23639800). Detected in erythrocytes (at protein level) (PubMed:11584005, PubMed:23219802). Expressed at relatively high levels in cerebral cortex, particularly the frontal and parietal lobes, and the thalamus and basal ganglia (at protein level) (PubMed:11311901). In the midbrain expressed at moderate levels, whereas in the other areas of the brainstem, namely medulla and pons, cerebellum and the hippocampus expressed at lower amounts when compared to the other brain regions (at protein level) (PubMed:11311901) Expressed in Langerhans cells and lymphocytes in the pancreas (at protein level) (PubMed:15501974). Expressed in kidney, in polarized renal epithelial cells (PubMed:12527552). Expressed in adipose tissues (PubMed:35790189). Expressed in placenta (PubMed:8986748). Expressed in small intestine (PubMed:10755314).

Background


Mediates both influx and efflux of nucleosides across the membrane (equilibrative transporter). It is sensitive (ES) to low concentrations of the inhibitor nitrobenzylmercaptapurine riboside (NBMPR) and is sodium-independent. It has a higher affinity for adenosine. Inhibited by dipyrindamole and dilazep (anticancer chemotherapeutics drugs).

References

- Griffiths M.,et al.Nat. Med. 3:89-93(1997).
Graham K.A.,et al.Submitted (SEP-1999) to the EMBL/GenBank/DDBJ databases.
Lum P.Y.,et al.Cancer Chemother. Pharmacol. 45:273-278(2000).
Sankar N.,et al.Nucleic Acids Res. 30:4339-4350(2002).
Mangravite L.M.,et al.Am. J. Physiol. 284:F902-F910(2003).

Images

Anti-ENT1 Antibodyat 1:1000 dilution + Hela whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L),Peroxidase conjugated at 1/10000 dilution Predicted band size : 50 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.