

GLUT1 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51519

Product Information

Application	WB
Primary Accession	P11166
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54084

Additional Information

Gene ID	6513
Other Names	Solute carrier family 2, facilitated glucose transporter member 1, Glucose transporter type 1, erythrocyte/brain, GLUT-1, HepG2 glucose transporter, SLC2A1, GLUT1
Dilution	WB~~1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	SLC2A1 (HGNC:11005)
Function	Facilitative glucose transporter, which is responsible for constitutive or basal glucose uptake (PubMed: 10227690 , PubMed: 10954735 , PubMed: 18245775 , PubMed: 19449892 , PubMed: 25982116 , PubMed: 27078104 , PubMed: 32860739). Has a very broad substrate specificity; can transport a wide range of aldoses including both pentoses and hexoses (PubMed: 18245775 , PubMed: 19449892). Most important energy carrier of the brain: present at the blood-brain barrier and assures the energy-independent, facilitative transport of glucose into the brain (PubMed: 10227690). In association with BSG and NXNL1, promotes retinal cone survival by increasing glucose uptake into photoreceptors (By similarity). Required for mesendoderm differentiation (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein. Melanosome. Photoreceptor inner segment {ECO:0000250 UniProtKB:P17809}. Note=Localizes primarily at the cell surface (PubMed:18245775, PubMed:19449892, PubMed:23219802, PubMed:24847886, PubMed:25982116). Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065)

Tissue Location	Detected in erythrocytes (at protein level). Expressed at variable levels in many human tissues
------------------------	---

Background

Facilitative glucose transporter. This isoform may be responsible for constitutive or basal glucose uptake. Has a very broad substrate specificity; can transport a wide range of aldoses including both pentoses and hexoses.

References

Mueckler M.,et al.Science 229:941-945(1985).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Fukumoto H.,et al.Diabetes 37:657-661(1988).
Yu W.,et al.Submitted (JUN-1998) to the EMBL/GenBank/DDBJ databases.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.