

Anti-GLUT3 Antibody

Catalog # AP51520

Product Information

Application	WB, IHC
Primary Accession	P11169
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53924

Additional Information

Gene ID	6515
Other Names	GLUT3; Solute carrier family 2, facilitated glucose transporter member 3; Glucose transporter type 3, brain; GLUT-3
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human GLUT3. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	SLC2A3 (HGNC:11007)
Function	Facilitative glucose transporter (PubMed: 26176916 , PubMed: 32860739 , PubMed: 9477959). Can also mediate the uptake of various other monosaccharides across the cell membrane (PubMed: 26176916 , PubMed: 9477959). Mediates the uptake of glucose, 2- deoxyglucose, galactose, mannose, xylose and fucose, and probably also dehydroascorbate (PubMed: 26176916 , PubMed: 9477959). Does not mediate fructose transport (PubMed: 26176916 , PubMed: 9477959). Required for mesendoderm differentiation (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein. Perikaryon {ECO:0000250 UniProtKB:Q07647}. Cell projection {ECO:0000250 UniProtKB:Q07647}. Note=Localized to densely spaced patches along neuronal processes. {ECO:0000250 UniProtKB:Q07647}
Tissue Location	Highly expressed in brain (PubMed:8457197). Expressed in many tissues.


References

Kayano T.,et al.J. Biol. Chem. 263:15245-15248(1988).
Stuart C.A.,et al.Submitted (JUN-2000) to the EMBL/GenBank/DDBJ databases.
Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

Images



Western blot analysis of GLUT3 expression in COLO205 (A), HepG2 (B) whole cell lysates. (Predicted band size: 53 kD; Observed band size: 53 kD)



Immunohistochemical analysis of GLUT3 staining in rat brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.