

Anti-STMN2 Antibody

Catalog # AP51541

Product Information

Application	WB
Primary Accession	Q93045
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Chicken, Bovine, Dog, Mk
Host	Rabbit
Clonality	Polyclonal
Calculated MW	20828

Additional Information

Gene ID	11075
Other Names	SCG10; SCGN10; Stathmin-2; Superior cervical ganglion-10 protein; Protein SCG10
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human STMN2. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000)
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	STMN2
Synonyms	SCG10, SCGN10
Function	Regulator of microtubule stability. When phosphorylated by MAPK8, stabilizes microtubules and consequently controls neurite length in cortical neurons. In the developing brain, negatively regulates the rate of exit from multipolar stage and retards radial migration from the ventricular zone (By similarity).
Cellular Location	Cytoplasm. Cytoplasm, perinuclear region. Cell projection, growth cone. Membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, axon. Golgi apparatus. Endosome. Cell projection, lamellipodium. Note=Associated with punctate structures in the perinuclear cytoplasm, axons, and growth cones of developing neurons. SCG10 exists in both soluble and membrane- bound forms. Colocalized with CIB1 in neurites of developing hippocampal primary neurons (By similarity). Colocalized with CIB1 in the cell

body, neuritis and growth cones of neurons. Colocalized with CIB1 to the leading edge of lamellipodia.


Tissue Location

Neuron specific.

References

Okazaki T.,et al.Neurobiol. Aging 16:883-894(1995).
Fujiwara T.,et al.Submitted (APR-1995) to the EMBL/GenBank/DDBJ databases.
Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).

Images



Western blot analysis of STMN2 expression in Jurkat (A), MCF7 (B), U2OS (C), mouse muscle (D) whole cell lysates. (Predicted band size: 20 kD; Observed band size: 25 kD)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.