

TCF3 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51551

Product Information

Application	WB, ICC, IHC-P
Primary Accession	<u>P15923</u>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67600

Additional Information

Gene ID	6929
Other Names	Transcription factor E2-alpha, Class B basic helix-loop-helix protein 21, bHLHb21, Immunoglobulin enhancer-binding factor E12/E47, Immunoglobulin transcription factor 1, Kappa-E2-binding factor, Transcription factor 3, TCF-3, Transcription factor ITF-1, TCF3, BHLHB21, E2A, ITF1
Dilution	WB~~1:1000 ICC~~N/A IHC-P~~N/A
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	TCF3
Synonyms	BHLHB21, E2A, ITF1
Function	Transcriptional regulator involved in the initiation of neuronal differentiation and mesenchymal to epithelial transition (By similarity). Heterodimers between TCF3 and tissue-specific basic helix- loop-helix (bHLH) proteins play major roles in determining tissue- specific cell fate during embryogenesis, like muscle or early B-cell differentiation (By similarity). Together with TCF15, required for the mesenchymal to epithelial transition (By similarity). Dimers bind DNA on E-box motifs: 5'-CANNTG-3' (By similarity). Binds to the kappa-E2 site in the kappa immunoglobulin gene enhancer (PubMed: <u>2493990</u>). Binds to IEB1 and IEB2, which are short DNA sequences in the insulin gene transcription control region (By similarity).
Cellular Location	Nucleus.

Background

Transcriptional regulator. Involved in the initiation of neuronal differentiation. Heterodimers between TCF3 and tissue- specific basic helix-loop-helix (bHLH) proteins play major roles in determining tissue-specific cell fate during embryogenesis, like muscle or early B-cell differentiation. Dimers bind DNA on E- box motifs: 5'-CANNTG-3'. Binds to the kappa-E2 site in the kappa immunoglobulin gene enhancer. Binds to IEB1 and IEB2, which are short DNA sequences in the insulin gene transcription control region.

References

Kamps M.P.,et al.Cell 60:547-555(1990). Nourse J.,et al.Cell 60:535-545(1990). Grimwood J.,et al.Nature 428:529-535(2004). Murre C.,et al.Cell 56:777-783(1989). Henthorn P.,et al.Nucleic Acids Res. 18:677-677(1990).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.