

# TGF beta 2 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51557

## Product Information

---

<b>Application</b>	WB, IHC-P
<b>Primary Accession</b>	<a href="#">P61812</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	47748

## Additional Information

---

<b>Gene ID</b>	7042
<b>Other Names</b>	Transforming growth factor beta-2, TGF-beta-2, BSC-1 cell growth inhibitor, Cetermin, Glioblastoma-derived T-cell suppressor factor, G-TSF, Polyergin, Latency-associated peptide, LAP, TGFB2
<b>Dilution</b>	WB~~1/500 - 1/1000 IHC-P~~N/A
<b>Format</b>	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
<b>Storage</b>	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

---

<b>Name</b>	TGFB2
<b>Function</b>	[Transforming growth factor beta-2 proprotein]: Precursor of the Latency-associated peptide (LAP) and Transforming growth factor beta-2 (TGF-beta-2) chains, which constitute the regulatory and active subunit of TGF-beta-2, respectively.
<b>Cellular Location</b>	[Latency-associated peptide]: Secreted, extracellular space, extracellular matrix {ECO:0000250   UniProtKB:P01137}

## Background

---

TGF-beta 2 has suppressive effects on interleukin-2 dependent T-cell growth.


## References

---


de Martin R.,et al.EMBO J. 6:3673-3677(1987).  
Madisen L.,et al.DNA 7:1-8(1988).  
Webb N.R.,et al.DNA 7:493-497(1988).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

## Images

---



Western blot analysis of TGF beta 2 expression in HEK293T (A); A375 (B); MCF7 (C) whole cell lysates.



Immunohistochemical analysis of TGF beta 2 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.