

# SCC-S2 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51572

## Product Information

---

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">O95379</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	23003

## Additional Information

---

<b>Gene ID</b>	25816
<b>Other Names</b>	Tumor necrosis factor alpha-induced protein 8, TNF alpha-induced protein 8, Head and neck tumor and metastasis-related protein, MDC-313, NF-kappa-B-inducible DED-containing protein, NDED, SCC-S2, TNF-induced protein GG2-1, TNFAIP8
<b>Target/Specificity</b>	KLH conjugated synthetic peptide derived from human SCC-S2
<b>Dilution</b>	WB~~ 1:1000
<b>Format</b>	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
<b>Storage</b>	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

---

<b>Name</b>	TNFAIP8
<b>Function</b>	Acts as a negative mediator of apoptosis and may play a role in tumor progression. Suppresses the TNF-mediated apoptosis by inhibiting caspase-8 activity but not the processing of procaspase-8, subsequently resulting in inhibition of BID cleavage and caspase-3 activation.
<b>Cellular Location</b>	Cytoplasm.
<b>Tissue Location</b>	Expressed at high levels in the spleen, lymph node, thymus, thyroid, bone marrow and placenta. Expressed at high levels both in various tumor tissues, unstimulated and cytokine-activated cultured cells. Expressed at low levels in the spinal cord, ovary, lung, adrenal glands, heart, brain, testis and skeletal muscle

## Background

---

Acts as a negative mediator of apoptosis and may play a role in tumor progression. Suppresses the TNF-mediated apoptosis by inhibiting caspase-8 activity but not the processing of procaspase-8, subsequently resulting in inhibition of BID cleavage and caspase-3 activation.

## References

---

Horrevoets A.J.G.,et al.Blood 93:3418-3431(1999).

Nietfeld W.,et al.Submitted (OCT-1998) to the EMBL/GenBank/DDBJ databases.


Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

## Images

---



Anti-SCC-S2 Antibody at 1:1000 dilution + human spleen lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 23 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.