10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-CD120b Antibody

Catalog # AP51574

Product Information

Application	WB, IHC, IF/IC
Primary Accession	P20333
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48291

Additional Information

Gene ID	7133
Other Names	TNFBR; TNFR2; Tumor necrosis factor receptor superfamily member 1B; Tumor necrosis factor receptor 2; TNF-R2; Tumor necrosis factor receptor type II; TNF-RII; TNFR-II; p75; p80 TNF-alpha receptor; CD120b; Etanercept
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human CD120b. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	TNFRSF1B
Synonyms	TNFBR, TNFR2
Function	Receptor with high affinity for TNFSF2/TNF and approximately 5-fold lower affinity for homotrimeric TNFSF1/lymphotoxin-alpha. The TRAF1/TRAF2 complex recruits the apoptotic suppressors BIRC2 and BIRC3 to TNFRSF1B/TNFR2. This receptor mediates most of the metabolic effects of TNF. Isoform 2 blocks TNF-induced apoptosis, which suggests that it regulates TNF function by antagonizing its biological activity.
Cellular Location	[Isoform 1]: Cell membrane; Single-pass type I membrane protein [Tumor necrosis factor-binding protein 2]: Secreted


References

Kohno T.,et al.Proc. Natl. Acad. Sci. U.S.A. 87:8331-8335(1990).
Smith C.A.,et al.Science 248:1019-1023(1990).
Beltinger C.P.,et al.Genomics 35:94-100(1996).
Lainez B.,et al.Int. Immunol. 16:169-177(2004).
Kalnine N.,et al.Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.


Images



Western blot analysis of CD120b expression in HEK293T (A), A375 (B), DLD (C), mouse brain (D), mouse heart (E), rat brain (F), rat heart (G) whole cell lysates. (Predicted band size: 48 kD; Observed band size: 65 kD)



Immunohistochemical analysis of CD120b staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of CD120b staining in PC12 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.