

14-3-3 gamma Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51619

Product Information

Application	WB
Primary Accession	P61981
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	28303

Additional Information

Gene ID	7532
Other Names	14-3-3 protein gamma, Protein kinase C inhibitor protein 1, KCIP-1, 14-3-3 protein gamma, N-terminally processed, YWHAG
Target/Specificity	KLH conjugated synthetic peptide derived from human 14-3-3 gamma
Dilution	WB~~ 1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	YWHAG (HGNC:12852)
Function	Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways (PubMed: 15696159 , PubMed: 16511572 , PubMed: 36732624). Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif (PubMed: 15696159 , PubMed: 16511572 , PubMed: 36732624). Binding generally results in the modulation of the activity of the binding partner (PubMed: 16511572). Promotes inactivation of WDR24 component of the GATOR2 complex by binding to phosphorylated WDR24 (PubMed: 36732624). Participates in the positive regulation of NMDA glutamate receptor activity by promoting the L- glutamate secretion through interaction with BEST1 (PubMed: 29121962). Reduces keratinocyte intercellular adhesion, via interacting with PKP1 and sequestering it in the cytoplasm, thereby reducing its incorporation into desmosomes (PubMed: 29678907). Plays a role in mitochondrial protein catabolic process (also named MALM) that promotes the degradation of damaged proteins inside mitochondria (PubMed: 22532927).

Cellular Location

Cytoplasm, cytosol. Mitochondrion matrix. Note=Translocates to the mitochondrial matrix following induction of MALM (mitochondrial protein catabolic process).

Tissue Location

Highly expressed in brain, skeletal muscle, and heart.

Background

Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways. Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif. Binding generally results in the modulation of the activity of the binding partner.

References

Autieri M.V.,et al.DNA Cell Biol. 18:555-564(1999).


Horie M.,et al.Genomics 60:241-243(1999).

Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.

Hillier L.W.,et al.Nature 424:157-164(2003).

Bienvenut W.V.,et al.Submitted (DEC-2008) to UniProtKB.

Images



All lanes : Anti-14-3-3 gamma Antibody at 1:1000 dilution
Lane 1: HeLa whole cell lysates Lane 2: NIH/3T3 whole cell lysates
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution
Predicted band size : 28 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.