

TTPAL Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP5164b

Product Information

Application	WB, IHC-P, FC, E
Primary Accession	Q9BTX7
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB26315
Calculated MW	38515
Antigen Region	204-233

Additional Information

Gene ID	79183
Other Names	Alpha-tocopherol transfer protein-like, TTPAL, C20orf121
Target/Specificity	This TTPAL antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 204-233 amino acids from the C-terminal region of human TTPAL.
Dilution	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TTPAL Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TTPAL
Synonyms	C20orf121
Function	May act as a protein that binds a hydrophobic ligand.


Background

TTPAL may act as a protein that binds a hydrophobic ligand.


References

Deloukas, P., et al. Nature 414(6866):865-871(2001)


Images



Western blot analysis of TTPAL Antibody (C-term) (Cat. #AP5164b) in MCF-7 cell line lysates (35ug/lane). TTPAL (arrow) was detected using the purified Pab.



TTPAL Antibody (C-term) (Cat. #AP5164b) IHC analysis in formalin fixed and paraffin embedded human breast carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the TTPAL Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



TTPAL Antibody (C-term) (Cat. #AP5164b) flow cytometric analysis of MCF-7 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.