

Anti-Kv1.3 (Phospho-Y187) Antibody

Catalog # AP51650

Product Information

Application	WB, IHC, IF/IC
Primary Accession	P22001
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Chicken, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	63842

Additional Information

Gene ID	3738
Other Names	HGK5; Potassium voltage-gated channel subfamily A member 3; HGK5; HLK3; HPCN3; Voltage-gated K(+) channel HuKIII; Voltage-gated potassium channel subunit Kv1.3
Target/Specificity	KLH-conjugated synthetic phosphopeptide corresponding to residues surrounding Y187 of human Kv1.3 protein. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	KCNA3
Synonyms	HGK5
Function	[Isoform 1]: Mediates the voltage-dependent potassium ion permeability of excitable membranes. Assuming opened or closed conformations in response to the voltage difference across the membrane, the protein forms a potassium-selective channel through which potassium ions may pass in accordance with their electrochemical gradient.
Cellular Location	[Isoform 1]: Cell membrane; Multi-pass membrane protein [Isoform 3]: Cytoplasm, perinuclear region


References

Folander K.,et al.Genomics 23:295-296(1994).
 Gregory S.G.,et al.Nature 441:315-321(2006).
 Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
 Philipson L.H.,et al.Proc. Natl. Acad. Sci. U.S.A. 88:53-57(1991).
 Attali B.,et al.J. Biol. Chem. 267:8650-8657(1992).


Images



Western blot analysis of Kv1.3 (Phospho-Y187) expression in HEK293T EGF-treated (A), SP2/0 EGF-treated (B), PC12 EGF-treated (C) whole cell lysates. (Predicted band size: 63 kD; Observed band size: 67 kD)



Immunohistochemical analysis of Kv1.3 (Phospho-Y187) staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of Kv1.3 (Phospho-Y187) staining in HuvEc cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.