

# TFE3 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51672

## Product Information

Application	WB, ICC
Primary Accession	<a href="#">P19532</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	61521

## Additional Information

Gene ID	7030
Other Names	Transcription factor E3, Class E basic helix-loop-helix protein 33, bHLHe33, TFE3, BHLHE33
Dilution	WB~~1:1000 ICC~~N/A
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

Name	TFE3 {ECO:0000303   PubMed:9393982, ECO:0000312   HGNC:HGNC:11752}
Function	<p>Transcription factor that acts as a master regulator of lysosomal biogenesis and immune response (PubMed:<a href="#">2338243</a>, PubMed:<a href="#">24448649</a>, PubMed:<a href="#">29146937</a>, PubMed:<a href="#">30733432</a>, PubMed:<a href="#">31672913</a>, PubMed:<a href="#">37079666</a>). Specifically recognizes and binds E-box sequences (5'-CANNTG-3'); efficient DNA-binding requires dimerization with itself or with another MiT/TFE family member such as TFEB or MITF (PubMed:<a href="#">24448649</a>). Involved in the cellular response to amino acid availability by acting downstream of MTOR: in the presence of nutrients, TFE3 phosphorylation by MTOR promotes its inactivation (PubMed:<a href="#">24448649</a>, PubMed:<a href="#">31672913</a>, PubMed:<a href="#">36608670</a>). Upon starvation or lysosomal stress, inhibition of MTOR induces TFE3 dephosphorylation, resulting in transcription factor activity (PubMed:<a href="#">24448649</a>, PubMed:<a href="#">31672913</a>, PubMed:<a href="#">36608670</a>). Specifically recognizes and binds the CLEAR-box sequence (5'-GTCACGTGAC-3') present in the regulatory region of many lysosomal genes, leading to activate their expression, thereby playing a central role in expression of lysosomal genes (PubMed:<a href="#">24448649</a>). Maintains the pluripotent state of embryonic stem cells by promoting the expression of genes such as ESRRB; mTOR- dependent TFE3 cytosolic retention and inactivation promotes exit from pluripotency (By</p>

similarity). Required to maintain the naive pluripotent state of hematopoietic stem cell; mTOR-dependent cytoplasmic retention of TFE3 promotes the exit of hematopoietic stem cell from pluripotency (PubMed:[30733432](#)). TFE3 activity is also involved in the inhibition of neuronal progenitor differentiation (By similarity). Acts as a positive regulator of browning of adipose tissue by promoting expression of target genes; mTOR-dependent phosphorylation promotes cytoplasmic retention of TFE3 and inhibits browning of adipose tissue (By similarity). In association with TFEB, activates the expression of CD40L in T-cells, thereby playing a role in T-cell- dependent antibody responses in activated CD4(+) T-cells and thymus- dependent humoral immunity (By similarity). Specifically recognizes the MUE3 box, a subset of E-boxes, present in the immunoglobulin enhancer (PubMed:[2338243](#)). It also binds very well to a USF/MLTF site (PubMed:[2338243](#)). Promotes TGF-beta-induced transcription of COL1A2; via its interaction with TSC22D1 at E-boxes in the gene proximal promoter (By similarity). May regulate lysosomal positioning in response to nutrient deprivation by promoting the expression of PIP4P1 (PubMed:[29146937](#)).

### Cellular Location

Cytoplasm, cytosol. Nucleus. Lysosome membrane. Note=When nutrients are present, recruited to the lysosomal membrane via association with GDP-bound RagC/RRAGC (or RagD/RRAGD); it is then phosphorylated by MTOR (PubMed:24448649, PubMed:37079666). Phosphorylation by MTOR prevents nuclear translocation and promotes ubiquitination and degradation (PubMed:22692423, PubMed:30733432, PubMed:36608670, PubMed:37079666) Conversely, inhibition of mTORC1, starvation and lysosomal disruption, promotes dephosphorylation and translocation to the nucleus (PubMed:22692423, PubMed:30733432, PubMed:37079666)

### Tissue Location

Ubiquitous in fetal and adult tissues.

## Background

Transcription factor that specifically recognizes and binds E-box sequences (5'-CANNTG-3'). Efficient DNA-binding requires dimerization with itself or with another MiT/TFE family member such as TFEB or MITF. In association with TFEB, activates the expression of CD40L in T-cells, thereby playing a role in T-cell-dependent antibody responses in activated CD4(+) T-cells and thymus-dependent humoral immunity. Specifically recognizes the MUE3 box, a subset of E-boxes, present in the immunoglobulin enhancer. It also binds very well to a USF/MLTF site.

## References

Clark J.,et al.Oncogene 15:2233-2239(1997).  
 Clark J.,et al.Submitted (NOV-1997) to the EMBL/GenBank/DDBJ databases.  
 Ross M.T.,et al.Nature 434:325-337(2005).  
 Weterman M.A.J.,et al.Proc. Natl. Acad. Sci. U.S.A. 93:15294-15298(1996).  
 Sidhar S.K.,et al.Hum. Mol. Genet. 5:1333-1338(1996).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.