

# Von Hippel Lindau Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51677

## Product Information

---

Application	WB
Primary Accession	<a href="#">P40337</a>
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	24153

## Additional Information

---

Gene ID	7428
Other Names	Von Hippel-Lindau disease tumor suppressor, Protein G7, pVHL, VHL
Dilution	WB~~1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

---

Name	VHL
Function	Involved in the ubiquitination and subsequent proteasomal degradation via the von Hippel-Lindau ubiquitination complex (PubMed: <a href="#">10944113</a> , PubMed: <a href="#">17981124</a> , PubMed: <a href="#">19584355</a> ). Seems to act as a target recruitment subunit in the E3 ubiquitin ligase complex and recruits hydroxylated hypoxia-inducible factor (HIF) under normoxic conditions (PubMed: <a href="#">10944113</a> , PubMed: <a href="#">17981124</a> ). Involved in transcriptional repression through interaction with HIF1A, HIF1AN and histone deacetylases (PubMed: <a href="#">10944113</a> , PubMed: <a href="#">17981124</a> ). Ubiquitinates, in an oxygen-responsive manner, ADRB2 (PubMed: <a href="#">19584355</a> ). Acts as a negative regulator of mTORC1 by promoting ubiquitination and degradation of RPTOR (PubMed: <a href="#">34290272</a> ).
Cellular Location	[Isoform 1]: Cytoplasm. Cell membrane; Peripheral membrane protein. Endoplasmic reticulum. Nucleus. Note=Found predominantly in the cytoplasm and with less amounts nuclear or membrane-associated (PubMed:9751722) Colocalizes with ADRB2 at the cell membrane (PubMed:19584355)
Tissue Location	Expressed in the adult and fetal brain and kidney.

## Background

---

Involved in the ubiquitination and subsequent proteasomal degradation via the von Hippel-Lindau ubiquitination complex. Seems to act as target recruitment subunit in the E3 ubiquitin ligase complex and recruits hydroxylated hypoxia- inducible factor (HIF) under normoxic conditions. Involved in transcriptional repression through interaction with HIF1A, HIF1AN and histone deacetylases. Ubiquitinates, in an oxygen-responsive manner, ADRB2.

## References

---

Latif F.,et al.Science 260:1317-1320(1993).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Muzny D.M.,et al.Nature 440:1194-1198(2006).  
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.  
Wenzel M.,et al.Submitted (APR-1996) to the EMBL/GenBank/DDBJ databases.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.