

# CD91 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51681

## Product Information

Application	WB
Primary Accession	<a href="#">Q07954</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	504606

## Additional Information

Gene ID	4035
Other Names	Prolow-density lipoprotein receptor-related protein 1, LRP-1, Alpha-2-macroglobulin receptor, A2MR, Apolipoprotein E receptor, APOER, CD91, Low-density lipoprotein receptor-related protein 1 85 kDa subunit, LRP-85, Low-density lipoprotein receptor-related protein 1 515 kDa subunit, LRP-515, Low-density lipoprotein receptor-related protein 1 intracellular domain, LRPICD, LRP1, A2MR, APR
Dilution	WB~~1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

Name	LRP1 ( <a href="#">HGNC:6692</a> )
Synonyms	A2MR, APR
Function	Endocytic receptor involved in endocytosis and in phagocytosis of apoptotic cells (PubMed: <a href="#">11907044</a> , PubMed: <a href="#">12713657</a> ). Required for early embryonic development (By similarity). Involved in cellular lipid homeostasis. Involved in the plasma clearance of chylomicron remnants and activated LRPAP1 (alpha 2-macroglobulin), as well as the local metabolism of complexes between plasminogen activators and their endogenous inhibitors. Acts as an LRPAP1 alpha-2- macroglobulin receptor (PubMed: <a href="#">1702392</a> , PubMed: <a href="#">26142438</a> ). Acts as TAU/MAPT receptor and controls the endocytosis of TAU/MAPT as well as its subsequent spread (PubMed: <a href="#">32296178</a> ). May modulate cellular events, such as APP metabolism, kinase-dependent intracellular signaling, neuronal calcium signaling as well as neurotransmission (PubMed: <a href="#">12888553</a> ). Also acts as a receptor for IGFBP3 to mediate cell growth inhibition (PubMed: <a href="#">9252371</a> ).

<b>Cellular Location</b>	[Low-density lipoprotein receptor-related protein 1 85 kDa subunit]: Cell membrane; Single-pass type I membrane protein Membrane, coated pit [Low-density lipoprotein receptor-related protein 1 intracellular domain]: Cytoplasm Nucleus. Note=After cleavage, the intracellular domain (LRP1CD) is detected both in the cytoplasm and in the nucleus.
<b>Tissue Location</b>	Most abundant in liver, brain and lung.

## Background

---

Endocytic receptor involved in endocytosis and in phagocytosis of apoptotic cells. Required for early embryonic development. Involved in cellular lipid homeostasis. Involved in the plasma clearance of chylomicron remnants and activated LRPAP1 (alpha 2-macroglobulin), as well as the local metabolism of complexes between plasminogen activators and their endogenous inhibitors. May modulate cellular events, such as APP metabolism, kinase-dependent intracellular signaling, neuronal calcium signaling as well as neurotransmission.

## References

---

Herz J., et al. EMBO J. 7:4119-4127(1988).  
 Van Leuven F., et al. Genomics 24:78-89(1994).  
 Van Leuven F., et al. Genomics 52:138-144(1998).  
 Scherer S.E., et al. Nature 440:346-351(2006).  
 Kutt H., et al. Biochim. Biophys. Acta 1009:229-236(1989).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.