

Anti-Cytochrome P450 2E1 Antibody

Catalog # AP51692

Product Information

Application	WB, IHC, IF/IC
Primary Accession	P05181
Reactivity	Human, Mouse, Rat, Rabbit, Dog, Mk
Host	Rabbit
Clonality	Polyclonal
Calculated MW	56849

Additional Information

Gene ID	1571
Other Names	CYP2E; Cytochrome P450 2E1; 4-nitrophenol 2-hydroxylase; CYP11E1; Cytochrome P450-J
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Cytochrome P450 2E1. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	CYP2E1 {ECO:0000303 PubMed:10553002, ECO:0000312 HGNC:HGNC:2631}
Function	A cytochrome P450 monooxygenase involved in the metabolism of fatty acids (PubMed: 10553002 , PubMed: 18577768). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (PubMed: 10553002 , PubMed: 18577768). Catalyzes the hydroxylation of carbon-hydrogen bonds. Hydroxylates fatty acids specifically at the omega-1 position displaying the highest catalytic activity for saturated fatty acids (PubMed: 10553002 , PubMed: 18577768). May be involved in the oxidative metabolism of xenobiotics (Probable).
Cellular Location	Endoplasmic reticulum membrane {ECO:0000250 UniProtKB:P05182}; Peripheral membrane protein {ECO:0000250 UniProtKB:P05182}. Microsome membrane {ECO:0000250 UniProtKB:P05182}; Peripheral membrane protein

{ECO:0000250|UniProtKB:P05182}. Mitochondrion inner membrane {ECO:0000250|UniProtKB:P05182}; Peripheral membrane protein {ECO:0000250|UniProtKB:P05182}. Note=Post-translationally targeted to mitochondria. TOMM70 is required for the translocation across the mitochondrial outer membrane. After translocation into the matrix, associates with the inner membrane as a membrane extrinsic protein {ECO:0000250|UniProtKB:P05182}


References

Song B.-J.,et al.J. Biol. Chem. 261:16689-16697(1986).
Umeno M.,et al.Biochemistry 27:9006-9013(1988).
Zhuge J.,et al.Submitted (SEP-1999) to the EMBL/GenBank/DDBJ databases.
Deloukas P.,et al.Nature 429:375-381(2004).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.


Images



Western blot analysis of Cytochrome P450 2E1 expression in Hela (A), U2OS (B), mouse liver (C), mouse brain (D), rat liver (E) whole cell lysates. (Predicted band size: 56 kD; Observed band size: 57 kD)



Immunohistochemical analysis of Cytochrome P450 2E1 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of Cytochrome P450 2E1 staining in NIH3T3 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.