

CLCN7 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51746

Product Information

Application WB, ICC **Primary Accession** WB, ICC

Reactivity Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Calculated MW 88679

Additional Information

Gene ID 1186

Other Names H(+)/Cl(-) exchange transporter 7, Chloride channel 7 alpha subunit, Chloride

channel protein 7, CIC-7, CLCN7

Dilution WB~~1:1000 ICC~~N/A

Format 0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name CLCN7 (HGNC:2025)

Function Slowly voltage-gated channel mediating the exchange of chloride ions

against protons (PubMed:<u>18449189</u>, PubMed:<u>21527911</u>). Functions as antiporter and contributes to the acidification of the lysosome lumen and

may be involved in maintaining lysosomal pH (PubMed: 18449189,

PubMed:<u>21527911</u>, PubMed:<u>31155284</u>). The CLC channel family contains both chloride channels and proton-coupled anion transporters that exchange chloride or another anion for protons (By similarity). The presence of conserved gating glutamate residues is typical for family members that

function as antiporters (By similarity).

Cellular Location Lysosome membrane; Multi-pass membrane protein

Tissue Location Brain and kidney...

Background

Slowly voltage-gated channel mediating the exchange of chloride ions against protons. Functions as antiporter and contributes to the acidification of the lysosome lumen.

References

Rae J.L.,et al.Submitted (JAN-2000) to the EMBL/GenBank/DDBJ databases. Ota T.,et al.Nat. Genet. 36:40-45(2004).

Martin J.,et al.Nature 432:988-994(2004).

Brandt S.,et al.FEBS Lett. 377:15-20(1995).

Eggermont J.,et al.Biochim. Biophys. Acta 1397:156-160(1998).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.