

# CLCNKA Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51747

## Product Information

Application	WB
Primary Accession	<a href="#">P51800</a>
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	75285

## Additional Information

Gene ID	1187
Other Names	Chloride channel protein CIC-Ka, Chloride channel Ka, CIC-K1, CLCNKA
Dilution	WB~~1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

Name	CLCNKA {ECO:0000303 PubMed:18310267, ECO:0000312 HGNC:HGNC:2026}
Function	Anion-selective channel permeable to small monovalent anions with ion selectivity for chloride > bromide > nitrate > iodide (PubMed: <a href="#">11734858</a> , PubMed: <a href="#">12111250</a> ). Forms a homodimeric channel where each subunit has its own ion conduction pathway. May conduct double- barreled currents controlled by two types of gates, two fast gates that control each subunit independently and a slow common gate that opens and shuts off both subunits simultaneously (PubMed: <a href="#">11734858</a> , PubMed: <a href="#">12111250</a> , PubMed: <a href="#">18310267</a> , PubMed: <a href="#">18776122</a> , PubMed: <a href="#">19646679</a> , PubMed: <a href="#">20538786</a> ). Assembles with the regulatory subunit BSND/Barttin for sorting at the basolateral plasma membrane domain and functional switch to the ion conducting state. CLCNKA:BSND channels display mostly a linear current-voltage relationship with fast gating at negative potentials (PubMed: <a href="#">11734858</a> , PubMed: <a href="#">12111250</a> , PubMed: <a href="#">18310267</a> , PubMed: <a href="#">18776122</a> , PubMed: <a href="#">20538786</a> ). Mediates transepithelial chloride transport from the lumen to interstitial compartment along the thin ascending limb of Henle's loop, contributing to generation of hypertonic medullary interstitium as a countercurrent system to achieve urine concentration (By similarity) (PubMed: <a href="#">15044642</a> ). Conducts chloride currents

in the stria vascularis of the inner ear to establish the endocochlear potential necessary for normal hearing (PubMed:[15044642](#), PubMed:[18310267](#), PubMed:[19646679](#)).

**Cellular Location**

Basolateral cell membrane {ECO:0000250|UniProtKB:Q9WUB7}; Multi-pass membrane protein

**Background**

---

Voltage-gated chloride channel. Chloride channels have several functions including the regulation of cell volume; membrane potential stabilization, signal transduction and transepithelial transport. May be important in urinary concentrating mechanisms.

**References**

---

Kieferle S.,et al.Proc. Natl. Acad. Sci. U.S.A. 91:6943-6947(1994).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Suzuki Y.,et al.Submitted (JUL-2006) to the EMBL/GenBank/DDBJ databases.  
Gregory S.G.,et al.Nature 441:315-321(2006).  
Schutte B.C.,et al.Submitted (MAR-1997) to the EMBL/GenBank/DDBJ databases.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.