

CD102 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51781

Product Information

Application	WB
Primary Accession	P13598
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	30654

Additional Information

Gene ID	3384
Other Names	Intercellular adhesion molecule 2, ICAM-2, CD102, ICAM2
Dilution	WB~~1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	ICAM2
Function	ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin alpha-L/beta-2). ICAM2 may play a role in lymphocyte recirculation by blocking LFA-1-dependent cell adhesion. It mediates adhesive interactions important for antigen-specific immune response, NK-cell mediated clearance, lymphocyte recirculation, and other cellular interactions important for immune response and surveillance.
Cellular Location	Membrane; Single-pass type I membrane protein. Cell projection, microvillus {ECO:0000250 UniProtKB:P35330}. Note=Co-localizes with RDX, EZR and MSN in microvilli. {ECO:0000250 UniProtKB:P35330}

Background

ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin alpha-L/beta-2). ICAM2 may play a role in lymphocyte recirculation by blocking LFA-1-dependent cell adhesion. It mediates adhesive interactions important for antigen- specific immune response, NK-cell mediated clearance, lymphocyte recirculation, and other cellular interactions important for immune response and surveillance.

References

Staunton D.E.,et al.Nature 339:61-64(1989).

Garcia-Aguilar J.,et al.Submitted (FEB-1990) to the EMBL/GenBank/DDBJ databases.

Xu G.,et al.Proc. Natl. Acad. Sci. U.S.A. 106:19310-19315(2009).

Liu T.,et al.J. Proteome Res. 4:2070-2080(2005).

Wollscheid B.,et al.Nat. Biotechnol. 27:378-386(2009).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.