

MPP9 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51793

Product Information

Application WB Primary Accession Q99550

Reactivity Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW133024

Additional Information

Gene ID 10198

Other Names M-phase phosphoprotein 9, MPHOSPH9, MPP9

Dilution WB~~1:1000

Format 0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name MPHOSPH9

Synonyms MPP9

Function Negatively regulates cilia formation by recruiting the CP110- CEP97 complex

(a negative regulator of ciliogenesis) at the distal end of the mother centriole in ciliary cells (PubMed:30375385). At the beginning of cilia formation,

MPHOSPH9 undergoes TTBK2-mediated phosphorylation and degradation via

the ubiquitin-proteasome system and removes itself and the CP110-CEP97 complex from the distal end of the mother centriole, which subsequently

promotes cilia formation (PubMed: 30375385).

Cellular Location Cytoplasm, cytoskeleton, microtubule organizing center, centrosome,

centriole. Golgi apparatus membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=Localizes to the distal and proximal end of centriole pairs in duplicated centrosomes. In ciliated cells, localizes to the distal and proximal end of daughter centriole and proximal of the mother centriole but not in the distal

end of the mother centriole (PubMed:21399614). Recruited by KIF24 to the

distal end of mother centriole where it forms a ring-like structure

(PubMed:30375385).

References

Scherer S.E.,et al.Nature 440:346-351(2006).

Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Matsumoto-Taniura N.,et al.Mol. Biol. Cell 7:1455-1469(1996).

Mayya V.,et al.Sci. Signal. 2:RA46-RA46(2009).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.