10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Kv1.5 Antibody

Catalog # AP51820

Product Information

Application	WB, IHC, IF/IC
Primary Accession	P22460
Reactivity	Human, Mouse, Rat, Pig, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67228

Additional Information

Gene ID	3741
Other Names	Potassium voltage-gated channel subfamily A member 5; HPCN1; Voltage-gated potassium channel HK2; Voltage-gated potassium channel subunit Kv1.5
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Kv1.5. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	KCNA5
Function	Voltage-gated potassium channel that mediates transmembrane potassium transport in excitable membranes. Forms tetrameric potassium- selective channels through which potassium ions pass in accordance with their electrochemical gradient. The channel alternates between opened and closed conformations in response to the voltage difference across the membrane. Can form functional homotetrameric channels and heterotetrameric channels that contain variable proportions of KCNA1, KCNA2, KCNA4, KCNA5, and possibly other family members as well; channel properties depend on the type of alpha subunits that are part of the channel (PubMed:12130714). Channel properties are modulated by cytoplasmic beta subunits that regulate the subcellular location of the alpha subunits and promote rapid inactivation (PubMed:12130714). Homotetrameric channels display rapid activation and slow inactivation (PubMed:12130714, PubMed:8505626). Required for normal electrical conduction including formation of the infranodal ventricular

	conduction system and normal action potential configuration, as a result of its interaction with XIRP2 (By similarity). May play a role in regulating the secretion of insulin in normal pancreatic islets.
Cellular Location	Cell membrane; Multi-pass membrane protein
Tissue Location	Pancreatic islets and insulinoma.


References

Tamkun M.M.,et al.FASEB J. 5:331-337(1991).
Philipson L.H.,et al.Proc. Natl. Acad. Sci. U.S.A. 88:53-57(1991).
Curran M.E.,et al.Genomics 12:729-737(1992).
Snyders D.J.,et al.J. Gen. Physiol. 101:513-543(1993).
Kurata H.T.,et al.J. Gen. Physiol. 118:315-332(2001).


Images



Western blot analysis of Kv1.5 expression in MCF7 (A), Raw264.7 (B), H9C2 (C) whole cell lysates. (Predicted band size: 67 kD; Observed band size: 56 kD)



Immunohistochemical analysis of Kv1.5 staining in human liver cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of Kv1.5 staining in RAW264.7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.